

Rabbit TATA Box Binding Protein Associated Factor 2 Polyclonal Antibody | anti-TAF2 antibody

TATA Box Binding Protein Associated Factor 2 (TAF2) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA145310	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

TATA Box Binding Protein Associated Factor 2; Polyclonal Antibody; TATA Box Binding Protein Associated Factor 2 (TAF2) Polyclonal Antibody; CIF150; TAF2B; TAFII150; TAF2 RNA Polymerase II; Transcription Initiation Factor TFIID Subunit 2; 150 kDa cofactor of initiator function RNA polymerase II TBP-associated factor B; anti-TAF2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	The antibody is a rabbit polyclonal antibody raised against TAF2. It has been selected for its ability to recognize TAF2 in immunohistochemical staining and western blotting.
Purity/Purification	Affinity Chromatography
Form/Format	Supplied as solution form in PBS, pH7.4 containing 0.02% NaN ₃ , 50% glycerol.
Concentration	500ug/mL (varies by lot)
Sequence Length	1199
Applicable Applications for anti-TAF2 antibody	ELISA, IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Immunoglobulin Type	IgG
Conjugate	No Conjugate
Cross Reactivity	Human
Organism Species	Human
Fragment	TAF2 (Pro918~Asp1199)
Quality Control	Content: The quality control contains recombinant TAF2 (Pro918~Asp1199) disposed in loading buffer. Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescient (ECL). Note: The quality control is specifically manufactured as the positive control. Not used for other purposes. Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN ₃ 0.02%.
Conjugated Antibody	The APC conjugated antibody version of this item is also available as
Preparation and Storage	Store at 4 degree C for frequent use. Stored at -20 degree C to -80 degree C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)
DAB staining on fromalin fixed paraffin- embedded Kidney tissue



WB (Western Blot)
Western Blot: Sample: Recombinant protein.



WB (Western Blot)
Western Blot: Sample: Human Hela cell lysate; Primary Ab: 1ug/ml Rabbit Anti-Human TAF2 Antibody Second Ab: 0.2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody



WB (Western Blot)
Western Blot: Sample: Human MCF7 cell lysate; Primary Ab: 1ug/ml Rabbit Anti-Human TAF2 Antibody Second Ab: 0.2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507347
NCBI GeneID	6873
NCBI Accession #	NP_003175.1
NCBI GenBank Nucleotide #	NM_003184.3
UniProt Accession #	Q6P1X5; O43487; O43604; O60668; Q86WW7; Q8IWK4; B2RE82
Molecular Weight	136,971 Da
NCBI Official Full Name	transcription initiation factor TFIID subunit 2
NCBI Official Synonym Full Names	TATA-box binding protein associated factor 2
NCBI Official Symbol	TAF2
NCBI Official Synonym Symbols	MRT40; TAF2B; CIF150; TAFII150
NCBI Protein Information	transcription initiation factor TFIID subunit 2
UniProt Protein Name	Transcription initiation factor TFIID subunit 2
UniProt Gene Name	TAF2
UniProt Synonym Gene Names	CIF150; TAF2B; TAF(II)150; TAFII-150; TAFII150

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tata-box-binding-protein-associated-factor-2-antibody-immunohistochemistry-immunocytochemistry-western-blot-elisa/145310>