

Vascular Endothelial Growth Factor A (VEGFA) Active Protein | VEGFA active protein

Active Vascular Endothelial Growth Factor A (VEGFA)

CATALOG # AAA146617	HOST 293F cell	APPLICATIONS Activity Assay, Cell Culture
-------------------------------	--------------------------	---


PRODUCT SYNONYMS

Vascular Endothelial Growth Factor A (VEGFA); Active Vascular Endothelial Growth Factor A (VEGFA); VEGF; VEGF-A; VPF; Vascular Endothelial Cell Growth Factor A; Vascular Permeability Factor; VEGFA active protein

PRODUCT SPECIFICATIONS

Host	293F cell
Purity/Purification	> 97%
Form/Format	Freeze-dried powder
Sequence Length	325
Applicable Applications for VEGFA active protein	Activity Assay, Cell Culture
Fragment	Ala27~Arg190
Tag	N-terminal His Tag
Buffer Formulation	PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Organism Species	Rat
Isoelectric Point	7.9
Usage	Reconstitute in PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
Preparation and Storage	Avoid repeated freeze/thaw cycles. Store at 2-8 degree C for one month. Aliquot and store at -80 degree C for 12 months. The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

IMAGES



Application Data






Figure 2. The dose-effect curve of VEGFA on ECV304 cells.


Application Data



Application Data



SDS-PAGE



Activity

Vascular endothelial growth factor A (VEGF-A), a glycosylated mitogen, is known to be a vascular permeability factor and an endothelial cell growth factor secreted by the smooth muscle and endothelial cells. It has been reported that VEGF-A induces vascular permeability and growth, promotes monocyte/macrophage migration, and inhibits cell apoptosis and so on. To test the effect of VEGF-A on cell proliferation of ECV304 endothelium cell line, cells were seeded into triplicate wells of 96-well plates at a density of 2,000 cells/well and allowed to attach overnight, then the medium was replaced with serum-free standard DMEM prior to the addition of various concentrations of VEGFA. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10uL of CCK-8 solution was added to each well of the plate, then measure the absorbance at 450nm using a microplate reader after incubating the plate for 1-4 hours at 37oC .Cell proliferation of ECV304 cells after incubation with VEGFA for 72h observed by inverted microscope was shown in Figure 1.

ADDITIONAL INFORMATION

Product Categories/Family for VEGFA active protein	Active Protein
--	----------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	160358797
NCBI GeneID	83785
NCBI Accession #	NP_001103804.1
NCBI GenBank Nucleotide #	NM_001110334.2
UniProt Accession #	P16612; Q541S6; Q91ZE1; Q9JKX7; Q9QXG6; Q9QXG7
Molecular Weight	20.8kDa; 22kDa
NCBI Official Full Name	vascular endothelial growth factor A isoform 3
NCBI Official Synonym Full Names	vascular endothelial growth factor A
NCBI Official Symbol	Vegfa
NCBI Official Synonym Symbols	VPF; Vegf; VEGF-A; VEGF111; VEGF164

NCBI Protein Information	vascular endothelial growth factor A
UniProt Protein Name	Vascular endothelial growth factor A
UniProt Gene Name	Vegfa
UniProt Synonym Gene Names	Vegf; VEGF-A; VPF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/293f-cell-active-protein/vascular-endothelial-growth-factor-a-vegfa/146617>