

Rat Hepcidin (Hepc) ELISA Kit | Hepc elisa kit

Rat Hepcidin (Hepc) ELISA Kit

CATALOG #
AAA147459

REACTIVITY
Rat


PRODUCT SYNONYMS

Hepcidin (Hepc); Rat Hepcidin (Hepc) ELISA Kit; HAMP; HFE2B; PLTR; LEAP1; Hepcidin Antimicrobial Peptide; Liver-expressed antimicrobial peptide 1; Putative liver tumor regressor; Hepc elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of Hepcidin (Hepc). No significant cross-reactivity or interference between Hepcidin (Hepc) and analogues was observed.
Sequence Length	84
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates And Other Biological Fluids
Detection Range	0.156-10ng/mL
Sensitivity	0.064ng/mL
Application	Enzyme-linked immunosorbent assay for Antigen Detection.
Intra-assay Precision (Precision within an assay)	3 samples with low, middle and high level Hepcidin (Hepc) were tested 20 times on one plate, respectively.
Inter-assay Precision (Precision between assays)	3 samples with low, middle and high level Hepcidin (Hepc) were tested on 3 different plates, 8 replicates in each plate.
CV(%) =	SD/meanX100
Intra-Assay	CV<10%
Inter-Assay	CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Hepc elisa kit	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Hepcidin (Hepc). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Hepcidin (Hepc). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Hepcidin (Hepc), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of Hepcidin (Hepc) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Product Categories/Family for Hepc elisa kit	Metabolic pathway; Endocrinology; Hematology; Genetic science

NCBI AND UNIPROT INFORMATION

NCBI GI #	16758218
NCBI GeneID	84604
NCBI Accession #	NP_445921.1
NCBI GenBank Nucleotide #	NM_053469.1
UniProt Accession #	Q99MH3
Molecular Weight	9,286 Da
NCBI Official Full Name	hepcidin
NCBI Official Synonym Full Names	hepcidin antimicrobial peptide
NCBI Official Symbol	Hamp
NCBI Official Synonym Symbols	Hepc
NCBI Protein Information	hepcidin
UniProt Protein Name	Hepcidin
UniProt Gene Name	Hamp
UniProt Synonym Gene Names	Hepc

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-elisa-kits/hepcidin-hepc/147459>