

Canine Creatine Kinase, Muscle (CKM) ELISA Kit | CKM elisa kit

Canine Creatine Kinase, Muscle (CKM) ELISA Kit

CATALOG #
AAA148979

REACTIVITY
Canine


PRODUCT SYNONYMS

Creatine Kinase; Muscle (CKM); Canine Creatine Kinase; Muscle (CKM) ELISA Kit; CKM elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Canine
Specificity	This assay has high sensitivity and excellent specificity for detection of CHUK. No significant cross-reactivity or interference between CHUK and analogues was observed.
Samples	Human Tissue Homogenates, Cell Lysates And Other Biological Fluids
Assay Type	Quantitative Sandwich
Detection Range	3.12-200pg/mL
Sensitivity	1.11pg/mL.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CHUK were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CHUK were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all the reagents should be kept according to the label on each vial. The Standard, Detection Reagent A, Detection Reagent B and the 96-well strip plate should be stored at -20 degree C while others should be at 4 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CKM elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CHUK in human tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to CHUK. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to CHUK. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CHUK, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CHUK in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/canine-elisa-kits/creatine-kinase-muscle-ckm/148979>