

Goat F(ab')₂ Anti-Human IgM (u chain specific) Secondary Antibody

Goat F(ab')₂ Anti-Human IgM-FITC


CATALOG # AAA14901	HOST Goat	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay
------------------------------	---------------------	----------------------------	---

PRODUCT SYNONYMS

Goat F(ab')₂ Anti-Human IgM (u chain specific); Polyclonal Secondary Antibody; Goat F(ab')₂ Anti-Human IgM-FITC; Goat F(ab')₂ Anti-Human IgM - FITC; Goat F(ab')₂ Anti-Human IgM (u chain specific) secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Reacts with the heavy chain of human IgM
Form/Format	Supplied as 0.5 mg in 1.0mL of PBS/NaN ₃ .
Applicable Applications for Goat F(ab') ₂ Anti-Human IgM (u chain specific) secondary antibody	FLISA, FCM/FACS (Flow Cytometry)
Application Notes	FLISA: FITC, AF488, and AF555 conjugates: 1:100-1:400 Flow Cytometry: FITC, BIOT, and AF488 conjugates: ≤1 ug/10⁶ cells Other applications: Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.
Source	Pepsin digest of goat anti-human IgM.
Cross Absorption	Human IgG and IgA; May reacts with IgM from other species.
Characterization	To ensure lot-to-lot consistency, each batch of product is tested by ELISA, Fluorescent ELISA, and/or flow cytometry for conformance to characteristics of a standard reference reagent.
Preparation and Storage	Store at 2-8 degree C. Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.



Application Data
Human peripheral blood lymphocytes were stained with Goat F(ab')₂ Anti-Human IgM-PE and Mouse Anti-Human CD19-FITC .

ADDITIONAL INFORMATION

Product Categories/Family for Goat F(ab') ₂ Anti-Human IgM (u chain specific) secondary antibody	Goat F (ab') ₂ Anti-Human IgM (u chain specific)
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.	

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/goat-fab2-anti-human-igm-u-chain-specific-secondary-antibody-flow-cytometry-elisa/14901>