

Rat Free Fatty Acid Receptor 3 (FFAR3) ELISA Kit | FFAR3 elisa kit

Rat Free Fatty Acid Receptor 3 (FFAR3) ELISA Kit

CATALOG #
AAA149022

REACTIVITY
Rat


PRODUCT SYNONYMS

Free Fatty Acid Receptor 3 (FFAR3); Rat Free Fatty Acid Receptor 3 (FFAR3) ELISA Kit; FFAR3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of FFAR2. No significant cross-reactivity or interference between FFAR2 and analogues was observed.
Sequence Length	346
Assay Type	Double-antibody Sandwich
Samples	Tissue homogenates, Cell lysates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	0.064ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level FFAR2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level FFAR2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FFAR3 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of FFAR2 in rat tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to FFAR2. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to FFAR2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FFAR2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of FFAR2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885329
NCBI GeneID	2865
NCBI Accession #	NP_005295.1
NCBI GenBank Nucleotide #	NM_005304.4
UniProt Accession #	O14843; Q14CM7; B2RWM8
Molecular Weight	38,649 Da
NCBI Official Full Name	free fatty acid receptor 3
NCBI Official Synonym Full Names	free fatty acid receptor 3
NCBI Official Symbol	FFAR3
NCBI Official Synonym Symbols	FFA3R; GPR41; GPR42
NCBI Protein Information	free fatty acid receptor 3
UniProt Protein Name	Free fatty acid receptor 3
UniProt Gene Name	FFAR3
UniProt Synonym Gene Names	GPR41

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-elisa-kits/free-fatty-acid-receptor-3-ffar3/149022>