

Goat F(ab')₂ Anti-Human IgG (gamma chain specific) Secondary Antibody

Goat F(ab')₂ Anti-Human IgG-FITC

CATALOG # AAA14907	HOST Goat	APPLICATIONS Flow Cytometry, Functional Assay
------------------------------	---------------------	---

PRODUCT SYNONYMS

Goat F(ab')₂ Anti-Human IgG (gamma chain specific); Polyclonal Secondary Antibody; Goat F(ab')₂ Anti-Human IgG-FITC; Goat F(ab')₂ Anti-Human IgG - FITC; Goat F(ab')₂ Anti-Human IgG (gamma chain specific) secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Specificity	Reacts with the heavy chain of human IgG
Form/Format	The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN ₃ .
Applicable Applications for Goat F(ab') ₂ Anti-Human IgG (gamma chain specific) secondary antibody	FLISA, FCM/FACS (Flow Cytometry)
Application Notes	FC: FITC ≤ 1 µg/10⁶ cells FLISA: 1:100-1:400 For flow cytometry, the suggested use of these reagents is in a final volume of 100 µl. Other Applications: Since applications vary, you should determine the optimum working dilution of the product that is appropriate for your specific need.
Source	Pepsin digest of goat anti-human IgG.
Cross Absorption	Human IgM, and IgA. May react with IgG from other species
Preparation and Storage	Store at 2-8 degrees C. Protect Fluorochrome-conjugated forms from light.

ADDITIONAL INFORMATION

Product Categories/Family for Goat F(ab') ₂ Anti-Human IgG (gamma chain specific) secondary antibody	Goat F(ab') ₂ Anti-Human IgG (? chain specific)
---	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/goat-fab2-anti-human-igg-gamma-chain-specific-secondary-antibody-flow-cytometry-elisa/14907>