

Goat F(ab')₂ Anti-Human IgM Secondary Antibody

Goat F(ab')₂ Anti-Human IgM-HRP

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA14912	Pepsin digest of Goat Anti-Human IgM	Human	ELISA, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Goat F(ab')₂ Anti-Human IgM; Goat F(ab')₂ Anti-Human IgM-HRP; Goat F(ab')₂ Anti-Human IgM (u chain specific); Goat F(ab')₂ Anti-Human IgM secondary antibody

PRODUCT SPECIFICATIONS

Host	Pepsin digest of Goat Anti-Human IgM
Reactivity	Human
Specificity	Reacts with the heavy chain of human IgM
Form/Format	Supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4
Applicable Applications for Goat F(ab') ₂ Anti-Human IgM secondary antibody	ELISA, FLISA, FCM/FACS (Flow Cytometry)
Application Notes	ELISA: 1:4,000 - 1:8,000 Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.
Cross Absorption	Human IgG and IgA; may react with IgM from other species.
Preparation and Storage	The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.

IMAGES



Application Data

Human peripheral blood lymphocytes were stained with Goat F(ab')₂ antihuman IgM-PE and mouse anti-human CD19-FITC.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.