

Rabbit Bcl2 Associated Athanogene 5 (BAG5) Polyclonal Antibody | anti-BAG5 antibody

Polyclonal Antibody to Bcl2 Associated Athanogene 5 (BAG5)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA149752	Rabbit	Rat, Human, Mouse	Immunoprecipitation, Immunocytochemistry, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Bcl2 Associated Athanogene 5 (BAG5); Polyclonal Antibody; Polyclonal Antibody to Bcl2 Associated Athanogene 5 (BAG5); BAG family molecular chaperone regulator 5; anti-BAG5 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Rat, Human, Mouse
Clonality	Polyclonal
Specificity	The antibody is a rabbit polyclonal antibody raised against BAG5. It has been selected for its ability to recognize BAG5 in immunohistochemical staining and western blotting.
Purity/Purification	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Form/Format	Liquid; PBS, pH7.4, containing 0.02% NaN ₃ , 50% glycerol.
Concentration	500ug/ml (varies by lot)
Applicable Applications for anti-BAG5 antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Cross Reactivity	Rat
Conjugation	None
Preparation and Storage	Store at 4 degree C for frequent use. Aliquot and store at -20 degree C for 12 months. Avoid repeated freeze/thaw cycles. The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

IMAGES



IHC (Immunohistochemistry)
DAB staining on formalin fixed paraffin-embedded kidney tissue



WB (Western Blot)
Western Blot

ADDITIONAL INFORMATION

Product Categories/Family for anti-BAG5 antibody	Tumor Immunity
--	----------------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

