

Cattle Thymic Factor (TF) ELISA Kit | TF elisa kit

Cattle Thymic Factor (TF) ELISA Kit

CATALOG #
AAA149896

REACTIVITY
Cattle


PRODUCT SYNONYMS

Thymic Factor (TF); Cattle Thymic Factor (TF) ELISA Kit; FTS; Facteur Thymique Serique; Thymulin; TF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Cattle
Specificity	This assay has high sensitivity and excellent specificity for detection of Thymic Factor (TF). No significant cross-reactivity or interference between Thymic Factor (TF) and analogues was observed.
Assay Type	Competitive Inhibition
Samples	Serum, Plasma and other biological fluids
Detection Range	37.0-3,000pg/mL
Sensitivity	13.3pg/mL
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Thymic Factor (TF) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Thymic Factor (TF) were tested on 3 different plates, 8 replicates in each plate. $CV(\%) = SD/mean \times 100$ Intra-Assay: $CV < 10\%$ Inter-Assay: $CV < 12\%$
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TF elisa kit	This assay employs the competitive inhibition enzyme immunoassay technique. A monoclonal antibody specific to Thymic Factor (TF) has been pre-coated onto a microplate. A competitive inhibition reaction is launched between biotin labeled Thymic Factor (TF) and unlabeled Thymic Factor (TF) (Standards or samples) with the pre-coated antibody specific to Thymic Factor (TF). After incubation the unbound conjugate is washed off. Next, avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. The amount of bound HRP conjugate is reverse proportional to the concentration of Thymic Factor (TF) in the sample. After addition of the substrate solution, the intensity of color developed is reverse proportional to the concentration of Thymic Factor (TF) in the sample.
Product Categories/Family for TF elisa kit	Infection Immunity; Endocrinology; Neuro Science; Hormone Metabolism

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/cat-cattle-elisa-kits/thymic-factor-tf/149896>