

General Oxidized Glutathione (GSSG) ELISA Kit | GSSG elisa kit

General Oxidized Glutathione (GSSG) ELISA Kit

CATALOG #
AAA149935

REACTIVITY
General


PRODUCT SYNONYMS

Oxidized Glutathione (GSSG); General Oxidized Glutathione (GSSG) ELISA Kit; GSSG elisa kit

PRODUCT SPECIFICATIONS

Reactivity	General
Specificity	This assay has high sensitivity and excellent specificity for detection of Oxidized Glutathione (GSSG). No significant cross-reactivity or interference between Oxidized Glutathione (GSSG) and analogues was observed.
Assay Type	Competitive
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates And Other Biological Fluids
Detection Range	246.9-20,000ng/mL
Sensitivity	88.4ng/mL
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Oxidized Glutathione (GSSG) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Oxidized Glutathione (GSSG) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for GSSG elisa kit	This assay employs the competitive inhibition enzyme immunoassay technique. A monoclonal antibody specific to Oxidized Glutathione (GSSG) has been pre-coated onto a microplate. A competitive inhibition reaction is launched between biotin labeled Oxidized Glutathione (GSSG) and unlabeled Oxidized Glutathione (GSSG) (Standards or samples) with the pre-coated antibody specific to Oxidized Glutathione (GSSG). After incubation the unbound conjugate is washed off. Next, avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. The amount of bound HRP conjugate is reverse proportional to the concentration of Oxidized Glutathione (GSSG) in the sample. After addition of the substrate solution, the intensity of color developed is reverse proportional to the concentration of Oxidized Glutathione (GSSG) in the sample.
--	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/general-elisa-kits/oxidized-glutathione-gssg/149935>