

Interleukin 6 Active Protein | IL6 active protein

Active Interleukin 6 (IL6)

CATALOG #
AAA150062

APPLICATIONS
Activity Assay, Cell Culture


PRODUCT SYNONYMS

Interleukin 6; Active Interleukin 6 (IL6); MGI2-A; MGI2A; HGF; BSF2; HSF; IFNB2; B-Cell Stimulatory Factor-2; Hybridoma/Plasmacytoma Growth Factor; Hepatocyte Stimulating Factor; Cytotoxic T-Cell Differentiation Factor; IL6 active protein

PRODUCT SPECIFICATIONS


Purity/Purification	>95%
Form/Format	Freeze-dried powder 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Sequence Positions	Ala18~Met207
Sequence Length	207
Applicable Applications for IL6 active protein	Activity Assay, Cell Culture
Species	Dog
Tag	N-terminal His Tag
Isoelectric Point	6.5
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
Preparation and Storage	Store at 2-8 degree C for one month. Aliquot and store at -80 degree C for 12 months. Avoid repeated freeze/thaw cycles. Stability: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

IMAGES




Application Data

Figure. Cell proliferation of Raji cells after stimulated with IL6. Interleukin 6 (IL-6) is an interleukin that acts as both a pro-inflammatory cytokine and an anti-inflammatory myokine. Interleukin 6 is secreted by T cells and macrophages to stimulate immune response and also plays a role in fighting infection. It supports the growth of B cells and is antagonistic to regulatory T cells. To test the effect of IL6 on cell proliferation, Raji cells were seeded into triplicate wells of 96-well plates at a density of 5,000 cells/well with 1% serum standard 1640 which contains various concentrations of recombinant dog IL6. After incubated for 5 days, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10uL of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37 degree C. Proliferation of Raji cells after incubation with IL6 for 5 days observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant IL6 for 5 days. The result was shown in Figure 2. It was obvious that IL6 significantly increased cell viability of Raji cells. (A) Raji cells cultured in 1640, stimulated with 1ng/mL IL6 for 5 days; (B) Unstimulated Raji cells cultured in 1640 for 5 days.



WB (Western Blot)
Figure. Western Blot



SDS-PAGE
Figure. SDS-PAGE

ADDITIONAL INFORMATION

Product Categories/Family for IL6 active protein	Cytokine; Tumor immunity; Infection immunity; Cardiovascular biology
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	729836
NCBI GeneID	403985
UniProt Accession #	P41323
Molecular Weight	Predicted Molecular Mass: 24.9kDa Accurate Molecular Mass: 25kDa
NCBI Official Full Name	Interleukin-6
NCBI Official Synonym Full Names	interleukin 6
NCBI Official Symbol	IL6
NCBI Official Synonym Symbols	IL-6

NCBI Protein Information	interleukin-6
UniProt Protein Name	Interleukin-6
UniProt Gene Name	IL6
UniProt Synonym Gene Names	IL-6

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/il6-active-protein/interleukin-6/150062>