

Pig Resistin (RETN) ELISA Kit | RETN elisa kit

Pig Resistin (RETN) ELISA Kit

CATALOG # AAA150455	REACTIVITY Pig	APPLICATIONS ELISA
-------------------------------	--------------------------	------------------------------

PRODUCT SYNONYMS

Resistin (RETN); Pig Resistin (RETN) ELISA Kit; ADSF; FIZZ3; RETN1; RSTN; XCP1; Adipose tissue-specific secretory factor; C/EBP-epsilon-regulated myeloid-specific secreted cysteine-rich protein; Cysteine-rich secreted FIZZ3; RETN elisa kit

PRODUCT SPECIFICATIONS


Reactivity	Pig
Specificity	This assay has high sensitivity and excellent specificity for detection of Resistin (RETN). No significant cross-reactivity or interference between Resistin (RETN) and analogues was observed.
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma, Tissue homogenates, Cell lysates, Cell culture supernates and other biological fluids
Detection Range	0.312-20ng/mL
Sensitivity	0.133ng/mL
Applicable Applications for RETN elisa kit	ELISA
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Resistin (RETN) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Resistin (RETN) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Resistin (RETN). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Resistin (RETN). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Resistin (RETN), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of Resistin (RETN) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Assay Procedure Summary	1. Prepare all reagents, samples and standards 2. Add 100uL standard or sample to each well. Incubate 2 hours at 37 degree C 3. Aspirate and add 100uL prepared Detection Reagent A. Incubate 1 hour at 37 degree C 4. Aspirate and wash 3 times 5. Add 100uL prepared Detection Reagent B. Incubate 30 minutes at 37 degree C 6. Aspirate and wash 5 times 7. Add 90uL Substrate Solution. Incubate 10-20 minutes at 37 degree C 8. Add 50uL Stop Solution. Read at 450nm immediately.

Preparation and Storage

The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.

To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Product Categories/Family for RETN elisa kit

Metabolic pathway; Endocrinology; Cardiovascular biology

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/pig-elisa-kits/resistin-retn/150455>