

Human Visfatin (VF) ELISA Kit | VF elisa kit

Human Visfatin (VF)(Sandwich ELISA) ELISA Kit

CATALOG # AAA150860	REACTIVITY Human	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------


PRODUCT SYNONYMS

Visfatin (VF); Human Visfatin (VF)(Sandwich ELISA) ELISA Kit; Mini Samples ELISA Kit for Visfatin (VF); VF/Visfatin; Nampt; NAmPRTase; PBEF; PBEF1; Pre-B-Cell Colony Enhancing Factor; Nicotinamide Phosphoribosyltransferase; VF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of visfatin. No significant cross-reactivity or interference between visfatin and analogues was observed.
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	1.56-100ng/mL
Sensitivity	< 0.59ng/mL
Applicable Applications for VF elisa kit	ELISA
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level visfatin were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level visfatin were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	Store at 4 degree C for 1 month; -20 degree C for long-term (within shelf life) The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for VF elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to visfatin. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to visfatin. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain visfatin, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of visfatin in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of visfatin in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Product Categories/Family for VF elisa kit	Metabolic pathway

NCBI AND UNIPROT INFORMATION

NCBI GI #	1172027
NCBI GeneID	10135
UniProt Accession #	P43490; Q3KQV0; Q8WW95; A4D0Q9; A4D0R0
Molecular Weight	491
NCBI Official Full Name	Nicotinamide phosphoribosyltransferase
NCBI Official Synonym Full Names	nicotinamide phosphoribosyltransferase
NCBI Official Symbol	NAMPT
NCBI Official Synonym Symbols	VF; PBEF; PBEF1; VISFATIN; 1110035O14Rik
NCBI Protein Information	nicotinamide phosphoribosyltransferase; NAMPRTase; pre-B cell-enhancing factor; pre-B-cell colony enhancing factor 1; pre-B-cell colony-enhancing factor 1
UniProt Protein Name	Nicotinamide phosphoribosyltransferase
UniProt Gene Name	NAMPT
UniProt Synonym Gene Names	PBEF; PBEF1; NAMPRTase; Nampt; Pre-B cell-enhancing factor
UniProt Entry Name	NAMPT_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.