

Rabbit anti-Pig Fibrinogen (FG) Polyclonal Antibody | anti-FG antibody

Polyclonal Antibody to Fibrinogen (FG)

CATALOG # AAA151919	HOST Rabbit	REACTIVITY Pig	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--------------------------	---

PRODUCT SYNONYMS

Fibrinogen (FG); Polyclonal Antibody; Polyclonal Antibody to Fibrinogen (FG); FG/Fibrinogen; F1; FI; Fbg; Coagulation Factor I; anti-FG antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Pig
Clonality	Polyclonal
Purity/Purification	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Form/Format	Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration	0.5mg/mL (varies by lot)
Applicable Applications for anti-FG antibody	IHC (Immunohistochemistry), WB (Western Blot)
Source	Polyclonal antibody preparation
Traits	Liquid
Cross Reactivity	Porcine
Immunogen	Native Fibrinogen .
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 4 degree C for frequent use. Aliquot and store at -20 degree C for 24 months. Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.




WB (Western Blot)

Western Blot; Samples: Lane 1: Porcine Liver lysate; Lane 2: Porcine Kidney lysate; Primary Ab: 0.01ug/ml Rabbit Anti-Porcine FG Antibody; Second Ab: 0.2ug/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody




IHC (Immunohistochemistry)

DAB staining on IHC-P; Sample: Porcine Kidney Tissue; Primary Ab: 10ug/ml Rabbit Anti-Porcine FG Antibody; Control: Used PBS instead of primary antibody; Second Ab: 2ug/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody




IHC (Immunohistochemistry)

DAB staining on IHC-P; Sample: Porcine Liver Tissue; Primary Ab: 10ug/ml Rabbit Anti-Porcine FG Antibody; Control: Used PBS instead of primary antibody; Second Ab: 2ug/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody




IHC (Immunohistochemistry)

DAB staining on IHC-P; Sample: Porcine Colon Tissue; Primary Ab: 10ug/ml Rabbit Anti-Porcine FG Antibody; Control: Used PBS instead of primary antibody; Second Ab: 2ug/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody



IHC (Immunohistochemistry)

DAB staining on IHC-P; Samples: Porcine Colon Tissue; Primary Ab: 10ug/ml Rabbit Anti-Porcine FG Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody



IHC (Immunohistochemistry)

DAB staining on IHC-P; Samples: Porcine Kidney Tissue; Primary Ab: 10ug/ml Rabbit Anti-Porcine FG Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

ADDITIONAL INFORMATION

Product Categories/Family for anti-FG antibody	Primary
--	---------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fibrinogen-fg-antibody-immunohistochemistry-western-blot-elisa/151919>