

Mouse Breakpoint Cluster Region (BCR) ELISA Kit | BCR elisa kit

Mouse Breakpoint Cluster Region (BCR)(Sandwich ELISA) ELISA Kit

CATALOG #
AAA152462

REACTIVITY
Mouse


PRODUCT SYNONYMS

Breakpoint Cluster Region (BCR); Mouse Breakpoint Cluster Region (BCR)(Sandwich ELISA) ELISA Kit; ELISA Kit for Breakpoint Cluster Region (BCR); ALL; BCR/FGFR1; BCR1; CML; FGFR1/BCR; PHL; Renal carcinoma antigen NY-REN-26; BCR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of BCR. No significant cross-reactivity or interference between BCR and analogues was observed.
Assay Type	Quantitative Sandwich
Samples	Mouse tissue homogenates, cell lysates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.061ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level BCR were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level BCR were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	Store at 4 degree C for 1 month/-20 degree C for long-term (within shelf life) Shelf-life: one year The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for BCR elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to BCR. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to BCR. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain BCR, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of BCR in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of BCR in mouse tissue homogenates, cell lysates and other biological fluids.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	124487229
NCBI GeneID	110279
NCBI Accession #	NP_001074881.1
NCBI GenBank Nucleotide #	NM_001081412.2
UniProt Accession #	Q6PAJ1; Q61339; Q6ZPE5; Q99LW5; E9PZL3
Molecular Weight	143,072 Da
NCBI Official Full Name	breakpoint cluster region protein
NCBI Official Synonym Full Names	breakpoint cluster region
NCBI Official Symbol	Bcr
NCBI Official Synonym Symbols	AI561783; AI853148; mKIAA3017; 5133400C09Rik
NCBI Protein Information	breakpoint cluster region protein; breakpoint cluster region homolog
UniProt Protein Name	Breakpoint cluster region protein
UniProt Gene Name	Bcr
UniProt Synonym Gene Names	Kiaa3017
UniProt Entry Name	BCR_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/breakpoint-cluster-region-bcr/152462>