

Mouse anti-Human Epidermal Growth Factor Receptor (EGFR) Monoclonal Antibody | anti-EGFR antibody

Monoclonal Antibody to Epidermal Growth Factor Receptor (EGFR)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA152603	Mouse	Human	Immunoprecipitation, Immunocytochemistry, Immunohistochemistry, Western Blot



PRODUCT SYNONYMS

Epidermal Growth Factor Receptor (EGFR); Monoclonal Antibody; Monoclonal Antibody to Epidermal Growth Factor Receptor (EGFR); Epidermal Growth Factor Receptor; ErbB-1; ErbB1; ER1; ERB-B1; MENA; HER1; Erythroblastic Leukemia Viral(v-Erb-B)oncogene Homolog; Avian; Proto-oncogene c-ErbB-1; Receptor tyrosine-protein kinase erbB-1; anti-EGFR antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG2b Kappa
Clone Number	C15
Specificity	The antibody is a mouse monoclonal antibody raised against EGFR. It has been selected for its ability to recognize EGFR in immunohistochemical staining and western blotting.
Purity/Purification	Affinity Chromatography
Form/Format	Liquid. 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration	1mg/mL (varies by lot)
Sequence Positions	His888~Ala1210
Applicable Applications for anti-EGFR antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant Epidermal Growth Factor Receptor (EGFR)
Preparation and Storage	Store at 4 degree C for frequent use. Store at -20 degree C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. Stable for 12 months if correctly stored. The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

IMAGES



IHC (Immunohistochemistry)
DAB staining on IHC-P; Sample: human Breast cancer Tissue; Primary Ab: 10ug/ml Mouse Anti-human EGFR Antibody Second Ab: 2ug/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody

WB (Western Blot)
Western Blot; Sample: Lane1: Hela cell lysate; Lane2: MCF7 cell lysate; Lane3: A431 cell lysate; Lane4: human Placenta lysate Primary Ab: 0.2ug/ml Mouse Anti-human EGFR Antibody Second Ab: 0.2ug/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody

ADDITIONAL INFORMATION

Product Categories/Family for anti-EGFR antibody	Monoclonal Antibodies (Primary Abs); Monoclonal antibodies; CD & Adhesion molecule
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	29725609
NCBI GeneID	1956
NCBI Accession #	NP_005219.2
NCBI GenBank Nucleotide #	NM_005228.3
UniProt Accession #	P00533; O00688; O00732; P06268; Q14225; Q68GS5; Q92795; Q9BZS2; Q9GZX1; Q9H2C9; Q9H3C9; Q9UMD7
Molecular Weight	134,277 Da
NCBI Official Full Name	epidermal growth factor receptor isoform a
NCBI Official Synonym Full Names	epidermal growth factor receptor
NCBI Official Symbol	EGFR
NCBI Official Synonym Symbols	ERBB; HER1; mENA; ERBB1; PIG61
NCBI Protein Information	epidermal growth factor receptor; proto-oncogene c-ErbB-1; cell growth inhibiting protein 40; cell proliferation-inducing protein 61; receptor tyrosine-protein kinase erbB-1; avian erythroblastic leukemia viral (v-erb-b) oncogene homolog
UniProt Protein Name	Epidermal growth factor receptor
UniProt Gene Name	EGFR
UniProt Synonym Gene Names	ERBB; ERBB1; HER1
UniProt Entry Name	EGFR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.