

Interleukin 6 (IL6) Active Protein | IL6 active protein

Active Interleukin 6 (IL6)



CATALOG # AAA161679	HOST 293F cell	APPLICATIONS Activity Assay, Cell Culture
-------------------------------	--------------------------	---

PRODUCT SYNONYMS

Interleukin 6 (IL6); Active Interleukin 6 (IL6); Interleukin 6; IL6 active protein (eukaryotic; 293F cell); Horse; MGI2-A; MGI2A; HGF; BSF2; HSF; IFNB2; B-Cell Stimulatory Factor-2; Hybridoma/Plasmacytoma Growth Factor; Hepatocyte Stimulating Factor; Cytotoxic T-Cell Differentiation Factor4; IL6 active protein

PRODUCT SPECIFICATIONS

Host	293F cell
Purity/Purification	Greater than 90% by SDS-PAGE
Form/Format	Freeze-dried powder; PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Sequence	Thr28~Met2 08
Applicable Applications for IL6 active protein	Activity Assay, Cell Culture
Species	Horse
Expression System	Eukaryotic expression
Tag	N-terminal His Tag
Endotoxin Level	<1.0EU per 1ug (determined by the LAL method)
Isoelectric Point	7.9
Research Area	Cytokine, Tumor immunity, Infection immunity, Cardiovascular biology
Preparation and Storage	Store at 2-8 degree C for one month, -80 degree C for 12 months



SDS-PAGE
Figure. SDS-PAGE

Bioactivity
Interleukin 6 (IL-6) is an interleukin that acts as both a pro-inflammatory cytokine and an anti-inflammatory myokine. Interleukin 6 is secreted by T cells and macrophages to stimulate immune response and also plays a role in fighting infection. It supports the growth of B cells and is antagonistic to regulatory T cells. To test the effect of IL6 on cell proliferation, TF-1 cells were seeded into triplicate wells of 96-well plates at a density of 20,000 cells/well with 10% serum standard 1640 which contains various concentrations of recombinant horse IL6. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 ul of CCK-8 solution was added to each well of the plate, then the absorbance at 450 nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37 degree C. Proliferation of TF-1 cells after incubation with IL6 for 72h observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant horse IL6 for 72h. The result was shown in Figure 2. It was obvious that IL6 significantly increased cell viability of TF-1 cells, the EC50 was 1.3 ng/ml.

ADDITIONAL INFORMATION

Product Categories/Family for IL6 active protein	Active proteins
---	-----------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	821159831
NCBI GeneID	100034196
NCBI Accession #	NP_001075965.2
NCBI GenBank Nucleotide #	NM_001082496.2
UniProt Accession #	Q95181
Molecular Weight	23,385 Da
NCBI Official Full Name	interleukin-6
NCBI Official Symbol	IL6
NCBI Official Synonym Symbols	IL-6
NCBI Protein Information	interleukin-6
UniProt Protein Name	Interleukin-6
UniProt Gene Name	IL6
UniProt Synonym Gene Names	IL-6

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/293f-cell-active-protein/interleukin-6-il6/161679>