

Oncostatin M (OSM) Active Protein | OSM active protein

Active Oncostatin M (OSM)


CATALOG # AAA161690	HOST 293F cell	APPLICATIONS Activity Assay, Cell Culture
-------------------------------	--------------------------	---

PRODUCT SYNONYMS

Oncostatin M (OSM); Active Oncostatin M (OSM); Oncostatin M; OSM active protein (eukaryotic; 293F cell); Human; OS-M4; OSM active protein

PRODUCT SPECIFICATIONS

Host	293F cell
Purity/Purification	Greater than 95% by SDS-PAGE
Form/Format	Freeze-dried powder; PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Sequence	Ala26~Arg2 20
Applicable Applications for OSM active protein	Activity Assay, Cell Culture
Species	Human
Expression System	Eukaryotic expression
Tag	N-terminal His Tag
Endotoxin Level	<1.0EU per 1ug (determined by the LAL method)
Isoelectric Point	9.4
Research Area	Cytokine, Tumor immunity
Preparation and Storage	Store at 2-8 degree C for one month, -80 degree C for 12 months




Bioactivity

Oncostatin M (OSM) is an approximately 30 KDa secreted cytokine belonging to the Interleukin-6 family. Like other members of the IL-6 family such as IL-11, CNTF, and Cardiotrophin-1, OSM plays crucial roles in inflammation, neuroprotection, hematopoiesis, metabolism and development. To test the effect of OSM on cell proliferation, TF-1 cells were seeded into triplicate wells of 96-well plates at a density of 20,000 cells/well with various concentrations of recombinant human OSM. After incubated for 72 hours, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8(CCK-8). Briefly, 10 ul of CCK-8 solution was added to each well of the plate, then the absorbance at 450 nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37 degree C. Proliferation of TF-1 cells after incubation with OSM for 72 hours observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with recombinant human OSM for 72 hours. The result was shown in Figure 2. It was obvious that recombinant human OSM significantly increased cell viability of TF-1 cells, the EC50 was 0.08 ng/ml.

SDS-PAGE

Figure. SDS-PAGE



ADDITIONAL INFORMATION

Product Categories/Family for OSM active protein	Active proteins
--	-----------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	10092621
NCBI GeneID	5008
NCBI Accession #	NP_065391.1
NCBI GenBank Nucleotide #	NM_020530.4
UniProt Accession #	P13725; Q6FHP8; Q9UCP6
Molecular Weight	28,484 Da
NCBI Official Full Name	oncostatin-M preproprotein
NCBI Official Synonym Full Names	oncostatin M
NCBI Official Symbol	OSM
NCBI Protein Information	oncostatin-M
UniProt Protein Name	Oncostatin-M
UniProt Gene Name	OSM
UniProt Synonym Gene Names	OSM
UniProt Entry Name	ONCM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/293f-cell-active-protein/oncostatin-m-osm/161690>