

Glutathione S Transferase Alpha 3 (GSTa3) Active Protein | GSTa3 active protein

Active Glutathione S Transferase Alpha 3 (GSTa3)

CATALOG # AAA161768	HOST E Coli	APPLICATIONS Activity Assay, Cell Culture
-------------------------------	-----------------------	---

PRODUCT SYNONYMS

Glutathione S Transferase Alpha 3 (GSTa3); Active Glutathione S Transferase Alpha 3 (GSTa3); Glutathione S Transferase Alpha 3; GST-A3; GSTA3-3; GTA3; GST class-alpha member 34; GSTa3 active protein

PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	Greater than 97% by SDS-PAGE
Form/Format	Freeze-dried powder; PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Sequence	Met1~Ser22 1
Applicable Applications for GSTa3 active protein	Activity Assay, Cell Culture
Species	Rat
Expression System	Prokaryotic expression
Tag	N-terminal His Tag
Endotoxin Level	<1.0EU per 1ug (determined by the LAL method)
Isoelectric Point	9.1
Research Area	Enzyme & Kinase, Tumor immunity, Hepatology
Preparation and Storage	Store at 2-8 degree C for one month, -80 degree C for 12 months

IMAGES

Specific Activity (pmol/min/ug)=

Adjusted V_{max} * (OD/min) x well volume (L) x 10^{12} pmol/mol
ext. coeff** (M⁻¹cm⁻¹) x path corr.*** (cm) x amount of enzyme (ug)


*Adjusted for Substrate Blank

**Using the extinction coefficient 9600 M⁻¹cm⁻¹

***Using the path correction 0.32 cm

Bioactivity

GSTa3 (Glutathione S-transferase a3) is an enzyme that plays an important role in detoxification by catalyzing the conjugation of many hydrophobic and electrophilic compounds with reduced glutathione. This subfamily of enzymes has a particular role in protecting cells from Reactive Oxygen Species and the products of peroxidation. Polymorphisms in this gene influence the ability of individuals to metabolize different drugs. GSTa3 catalyze the endogenous glutathione conjugation 1-Chloro-2,4-dinitrobenzene (CDNB), which can increase in the absorbance at 340 nm. The reaction was performed in adding 10 ul 200 mM glutathione (reduced) and 10 ul 100 mM CDNB in 980 ul 100 mM NaH₂PO₄ (pH7.0), rapidly mixed. Then add 50 ul mixed substrates to 50 ul different concentrations of recombinant rat GSTa3, mix gently. Incubated at 37 degree C for 5min, then read at a wavelength of 340 nm. The specific activity of recombinant rat GSTa3 is >37000 pmol/min/ug.



SDS-PAGE

Figure. SDS-PAGE

ADDITIONAL INFORMATION

Product Categories/Family for GSTa3 active protein	Active proteins
--	-----------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	13928688
NCBI GeneID	24421
NCBI Accession #	NP_113697.1
NCBI GenBank Nucleotide #	NM_031509.2
UniProt Accession #	P04904; Q6LD92
Molecular Weight	25,319 Da
NCBI Official Full Name	glutathione S-transferase alpha-3
NCBI Official Synonym Full Names	glutathione S-transferase alpha 1
NCBI Official Symbol	Gsta1
NCBI Official Synonym Symbols	Yc1; Gsta3; Gsta5; GST AA; GST 2-2; GST Yc1; GST A3-3
NCBI Protein Information	glutathione S-transferase alpha-3
UniProt Protein Name	Glutathione S-transferase alpha-3
UniProt Gene Name	Gsta3
UniProt Synonym Gene Names	Gstyc1; GST Yc1
UniProt Entry Name	GSTA3_RAT

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/e-coli-active-protein/glutathione-s-transferase-alpha-3-gsta3/161768>