

Apolipoprotein A5 (APOA5) Active Protein | APOA5 active protein

Active Apolipoprotein A5 (APOA5)


CATALOG # AAA161872	HOST E Coli	APPLICATIONS Activity Assay, Cell Culture
-------------------------------	-----------------------	---

PRODUCT SYNONYMS

Apolipoprotein A5 (APOA5); Active Apolipoprotein A5 (APOA5); Apolipoprotein A5; Apo-A5; APOA-V; APOAV; RAP3; Regeneration-associated protein 34; APOA5 active protein

PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	Greater than 90% by SDS-PAGE
Form/Format	Freeze-dried powder; PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Sequence	Asp167~Lys 335
Applicable Applications for APOA5 active protein	Activity Assay, Cell Culture
Species	Human
Expression System	Prokaryotic expression
Tag	N-terminal His Tag
Endotoxin Level	<1.0EU per 1ug (determined by the LAL method)
Isoelectric Point	6.7
Research Area	Metabolic pathway, Endocrinology, Cardiovascular biology, Hepatology, Nutrition metabolism
Preparation and Storage	Store at 2-8 degree C for one month, -80 degree C for 12 months




Bioactivity

Apolipoprotein A5 (APOA5) is a small protein, expressed predominantly in the liver. It has been identified to play an important role in lipid metabolism, specifically in triglyceride (TG) and TG-rich lipoproteins (TRLs) metabolism. Apolipoprotein C3 (APOC3) has been proved to be one of the ligands of APOA5. Thus a functional binding ELISA assay was conducted to detect the interaction of recombinant human APOA5 and recombinant human APOC3. Briefly, APOA5 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 ul were then transferred to APOC3-coated microtiter wells and incubated for 1h at 37 degree C. Wells were washed with PBST and incubated for 1h with anti-APOA5 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody for 1h at 37 degree C, wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37 degree C. Finally, add 50 uL stop solution to the wells and read at 450/630 nm immediately. The binding activity of recombinant human APOA5 and recombinant human APOC3 was shown in Figure 1, the EC50 for this effect is 0.14 ug/mL.

SDS-PAGE

Figure. SDS-PAGE



ADDITIONAL INFORMATION

Product Categories/Family for APOA5 active protein	Active proteins
--	-----------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	262231737
NCBI GeneID	116519
NCBI Accession #	NP_001160070.1
NCBI GenBank Nucleotide #	NM_001166598.1
UniProt Accession #	Q6Q788; Q3MIK6; Q6UWK9; Q9UBJ3; B0YIV9
Molecular Weight	41,213 Da
NCBI Official Full Name	apolipoprotein A-V
NCBI Official Synonym Full Names	apolipoprotein A-V
NCBI Official Symbol	APOA5
NCBI Official Synonym Symbols	RAP3; APOAV
NCBI Protein Information	apolipoprotein A-V; apo-AV; apolipoprotein A5; regeneration-associated protein 3
UniProt Protein Name	Apolipoprotein A-V
UniProt Gene Name	APOA5
UniProt Synonym Gene Names	RAP3; Apo-AV; ApoA-V
UniProt Entry Name	APOA5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/e-coli-active-protein/apolipoprotein-a5-apoa5/161872>