

Rabbit anti-Mouse, Human CCL17/TARC Polyclonal Antibody | anti-CCL17 antibody

CCL17/TARC Rabbit anti-Human Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA162910	Rabbit	Mouse, Human	Western Blot, Immunohistochemistry, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

CCL17/TARC; Polyclonal Antibody; CCL17/TARC Rabbit anti-Human Polyclonal Antibody; IHC-plus CCL17/TARC Antibody; CCL17; A-152E5.3; C-C motif chemokine 17; CC chemokine abcd-2; ABCD-2; SCYA17; T cell-directed CC chemokine; TARC; CC chemokine TARC; Small-inducible cytokine A17; anti-CCL17 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse, Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Human CCL17/TARC
Purity/Purification	Affinity purification
Form/Format	PBS, pH7.4, 0.05% Sodium Azide, 50% Glycerol
Concentration	0.8mg/ml (varies by lot)
Applicable Applications for anti-CCL17 antibody	WB (Western Blot), IHC (Immunohistochemistry), IHC (Immunohistochemistry), ELISA
Target	Human CCL17/TARC
Immunogen	Synthetic peptide of human CCL17
Conjugation	Unconjugated
Family	Interocrine
Subfamily	Interocrine beta
Preparation and Storage	Store at -20 degree C. Avoid freeze-thaw cycles.


IMAGES




IHC (Immunohistochemistry)
CCL17/TARC Antibody-Immunohistochemistry of paraffin-embedded Human colon cancer using CCL17 Polyclonal Antibody at dilution of 1:70.



IHC (Immunohistochemistry)
CCL17/TARC Antibody-Immunohistochemistry of paraffin-embedded Human gastric cancer using CCL17 Polyclonal Antibody at dilution of 1:70.



IHC (Immunohistochemistry)
CCL17/TARC Antibody-Human Spleen: Formalin-Fixed, Paraffin-Embedded (FFPE), at a dilution of 1:100.



WB (Western Blot)
CCL17/TARC Antibody-Western blot analysis of Mouse lung and thymus tissue, human fetal muscle tissue, using CCL17 Polyclonal Antibody at dilution of 1:1300.

ADDITIONAL INFORMATION

Related Product Information for anti-CCL17 antibody	TARC antibody is an unconjugated rabbit polyclonal antibody to TARC (CCL17) from human. It is reactive with human and mouse. Validated for ELISA, IHC and WB. Tested on 20 paraffin-embedded human tissues.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506829
NCBI GeneID	6361
NCBI Accession #	NP_002978.1
NCBI GenBank Nucleotide #	NM_002987
UniProt Accession #	Q92583; Q2M287; A0N0Q9
Molecular Weight	10,507 Da
NCBI Official Full Name	C-C motif chemokine 17
NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 17
NCBI Official Symbol	CCL17
NCBI Official Synonym Symbols	TARC; ABCD-2; SCYA17; A-152E5.3
NCBI Protein Information	C-C motif chemokine 17; CC chemokine TARC; T cell-directed CC chemokine; small inducible cytokine subfamily A (Cys-Cys), member 17; small-inducible cytokine A17; thymus and activation-regulated chemokine

UniProt Protein Name	C-C motif chemokine 17
UniProt Gene Name	CCL17
UniProt Synonym Gene Names	SCYA17; TARC
UniProt Entry Name	CCL17_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ccl17-tarc-antibody-western-blot-immunohistochemistry-immunohistochemistry-elisa/162910>