

Rabbit anti-Human KRT20/CK20/Cytokeratin 20 Polyclonal Antibody | anti-KRT20 antibody

KRT20/CK20/Cytokeratin 20 Rabbit anti-Human Polyclonal Antibody

CATALOG # AAA162957	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

KRT20/CK20/Cytokeratin 20; Polyclonal Antibody; KRT20/CK20/Cytokeratin 20 Rabbit anti-Human Polyclonal Antibody; PathPlus KRT20/CK20/Cytokeratin 20 Antibody; KRT20; Cytokeratin-20; Cytokeratin 20; CK20; K20; Keratin 20; Keratin-20; KRT21; CK-20; Protein IT; anti-KRT20 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Immunoaffinity purified
Form/Format	PBS, 0.1% Sodium Azide
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-KRT20 antibody	IHC (Immunohistochemistry)
Target	Human KRT20/CK20/Cytokeratin 20
Immunogen	LP-10498, synthetic peptide
Conjugation	Unconjugated
Family	Keratin
Preparation and Storage	Aliquot and store undiluted at -20 degree C or below for up to 1 year. Can be stored undiluted at 4 degree C for up to 1 month. Avoid freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

KRT20/CK20/Cytokeratin 20 Antibody-Human Small Intestine: Formalin-Fixed, Paraffin-Embedded (FFPE)



IHC (Immunohistochemistry)

KRT20/CK20/Cytokeratin 20 Antibody-Human Breast Carcinoma: Formalin-Fixed, Paraffin-Embedded (FFPE)



IHC (Immunohistochemistry)
KRT20/CK20/Cytokeratin 20 Antibody-Human Colon Carcinoma: Formalin-Fixed, Paraffin-Embedded (FFPE)

ADDITIONAL INFORMATION

Related Product Information for anti-KRT20 antibody	Cytokeratin 20 antibody is an unconjugated rabbit polyclonal antibody to Cytokeratin 20 (KRT20/CK20) from human. Validated for IHC. CK20 (KRT20) is an intermediate filament keratin that is expressed in epithelial cells in enterocytes and goblet cells of the gastrointestinal mucosa, in some endocrine cells and Merkel cells of the epidermis. It is frequently used as a differentiation marker in the colon, and is used in conjunction with CK7 to distinguish colon carcinomas (CK20 +, CK7 -) from ovarian, pulmonary, or breast cancers. Poorly differentiated colorectal tumors and those with high rates of MSI may also stain negatively for this marker. It is also positive in adenocarcinomas of the gallbladder and bile ducts, ductal cell adenocarcinomas of the pancreas, cholangiocarcinomas, mucinous ovarian tumors, transitional-cell carcinomas, and Merkel-cell carcinomas of the skin. CK20 is usually negative in squamous cell carcinomas, breast cancers, lung adenocarcinomas, endometrial cancer, and small cell carcinomas. Staining for this target is expected to be cytoplasmic.
References	Proc Natl Acad Sci U S A. 2009 Feb 10;106(6):1936-41, PMID: 19188603; Histology and Histopathology 27 (3):347-56. 2012/03, PMID: 22237712; Cell Stem Cell 8 (5):511-24. 5/6/2011, DOI: 10.1016/j.stem.2011.02.020;

NCBI AND UNIPROT INFORMATION

NCBI GI #	406854
UniProt Accession #	P35900; X73502
Molecular Weight	48,487 Da
NCBI Official Full Name	cytokeratin 20
UniProt Protein Name	Keratin, type I cytoskeletal 20
UniProt Gene Name	KRT20
UniProt Synonym Gene Names	CK-20; K20
UniProt Entry Name	K1C20_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/krt20-ck20-cytokeratin-20-antibody-immunohistochemistry-elisa/162957>