

Porcine E-Cad ELISA Kit | E-Cad elisa kit

Porcine E-Cad (E-Cadherin) ELISA Kit

CATALOG #
AAA164284

REACTIVITY
Porcine


PRODUCT SYNONYMS

E-Cad; Porcine E-Cad (E-Cadherin) ELISA Kit; E-Cad elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Porcine
Specificity	This kit recognizes Porcine E-Cad in samples. No significant cross-reactivity or interference between Porcine E-Cad and analogues was observed
Sequence Length	895
Assay Type	Quantitative Sandwich
Samples	Serum, plasma and other biological fluids
Detection Range	0.16-10 ng/mL
Sensitivity	0.09 ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, mid range and high level Porcine E-Cad were tested 20 times on one plate, respectively.
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, mid range and high level Porcine E-Cad were tested on 3 different plates, 20 replicates in each plate, respectively.
Preparation and Storage	Store at 4 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for E-Cad elisa kit	Principle of the Assay: This ELISA kit uses the Sandwich-ELISA principle. The micro ELISA plate provided in this kit has been pre-coated with an antibody specific to Porcine E-Cad. Samples (or Standards) are added to the micro ELISA plate wells and combined with the specific antibody. Then a biotinylated detection antibody specific for Porcine E-Cad and Avidin-Horseradish Peroxidase (HRP) conjugate are added successively to each micro plate well and incubated. Free components are washed away. The substrate solution is added to each well. Only those wells that contain Porcine E-Cad, biotinylated detection antibody and Avidin-HRP conjugate will appear blue in color. The enzyme-substrate reaction is terminated by the addition of stop solution and the color turns yellow. The optical density (OD) is measured spectrophotometrically at a wavelength of 450 ± 2 nm. The OD value is proportional to the concentration of Porcine E-Cad. You can calculate the concentration of Porcine E-Cad in the samples by comparing the OD of the samples to the standard curve. Intended Uses: This ELISA kit applies to the in vitro quantitative determination of Porcine E-Cad concentrations in serum, plasma and other biological fluids.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	148236899
NCBI GeneID	394290
NCBI Accession #	NP_001080946.1
NCBI GenBank Nucleotide #	NM_001087477.1
UniProt Accession #	P33148; Q91543
Molecular Weight	97,651 Da
NCBI Official Full Name	EP-cadherin
NCBI Official Synonym Full Names	cadherin 3, type 1, P-cadherin (placental)<
NCBI Official Symbol	cdh3-a
NCBI Official Synonym Symbols	uvo; cdhp; hjmd; lcam; pcad; E-Cad; XBcad; l-cam; xcdh1; cdh1-a; cdhc-A; xb-cad; XTCAD-1; E-cadherin; U-cadherin; b-cadherin; cadherin-1; uvomorulin; XB-cadherin
NCBI Protein Information	EP-cadherin; C-cadherin (EP-cadherin); cadherin 1, type 1, E-cadherin (epithelial)
UniProt Protein Name	EP-cadherin
UniProt Entry Name	CADHF_XENLA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.