

Human CgA ELISA Kit | CgA/Pancreastatin elisa kit

Human CgA (Chromogranin A/Pancreastatin) ELISA Kit

CATALOG #
AAA165420

REACTIVITY
Human


PRODUCT SYNONYMS

CgA; Human CgA (Chromogranin A/Pancreastatin) ELISA Kit; CgA/Pancreastatin elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of Human CgA/Pancreastatin. No significant cross-reactivity or interference between Human CgA/Pancreastatin and analogues was observed. Note: Limited by current skills and knowledge, it is impossible for us to complete the cross- reactivity detection between Human CgA/Pancreastatin and all the analogues, therefore, cross reaction may still exist.
Sequence Length	116
Samples	Serum, Plasma, Biological Fluids
Assay Type	Sandwich
Detection Range	62.5-4000pg/mL
Sensitivity	Min: 37.5pg/mL; Max: 4000pg/mL
Preparation and Storage	Store at 4 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CgA/Pancreastatin elisa kit	Description: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CgA/Pancreastatin in human serum, plasma and other biological fluids Principle of the Assay: This ELISA kit uses Sandwich-ELISA as the method. The micro ELISA plate provided in this kit has been pre-coated with an antibody specific to Human CgA/Pancreastatin. Standards or samples are added to the appropriate micro ELISA plate wells and bound by the specific antibody. Then a biotinylated detection antibody specific for Human CgA/Pancreastatin and Avidin-Horseradish Peroxidase (HRP) conjugate is added to each micro plate well successively and incubated. Free components are washed away. The substrate solution is added to each well. Only those wells that contain Human CgA/Pancreastatin, biotinylated detection antibody and Avidin-HRP conjugate will appear blue in color. The enzyme-substrate reaction is terminated by the addition of a sulphuric acid solution and the color turns yellow. The optical density (OD) is measured spectrophotometrically at a wavelength of 450 nm +/- 2 nm. The OD value is proportional to the concentration of Human CgA/Pancreastatin. You can calculate the concentration of Human CgA/Pancreastatin in the samples by comparing the OD of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	48146971
NCBI GeneID	1081
UniProt Accession #	Q6I9S8; P01215
Molecular Weight	13,075 Da
NCBI Official Full Name	CGA
NCBI Official Synonym Full Names	glycoprotein hormones, alpha polypeptide
NCBI Official Symbol	CGA
NCBI Official Synonym Symbols	HCG; LHA; FSHA; GPHa; TSHA; GPHA1; CG-ALPHA
NCBI Protein Information	glycoprotein hormones alpha chain; FSH-alpha; LSH-alpha; TSH-alpha; anterior pituitary glycoprotein hormones common subunit alpha; choriogonadotropin alpha chain; chorionic gonadotrophin subunit alpha; chorionic gonadotropin, alpha polypeptide; follicle-stimulating hormone alpha chain; follicle-stimulating hormone alpha subunit; follitropin alpha chain; luteinizing hormone alpha chain; lutropin alpha chain; thyroid-stimulating hormone alpha chain; thyrotropin alpha chain
UniProt Protein Name	CGA protein
UniProt Gene Name	CGA
UniProt Entry Name	Q6I9S8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/cga-pancreastatin-human-elisa-kits/cga/165420>