

Porcine Prolactin ELISA Kit | PRL elisa kit

Porcine Prolactin ELISA Kit

CATALOG #
AAA16583

REACTIVITY
Porcine


PRODUCT SYNONYMS

Prolactin; Porcine Prolactin ELISA Kit; PRL elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Porcine
Specificity	This assay has high sensitivity and excellent specificity for detection of PRL. No significant cross-reactivity or interference between PRL and analogues was observed. NOTE: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between PRL and all the analogues, therefore, cross reaction may still exist in some cases.
Samples	Serum, plasma, cell culture supernatants, body fluid and tissue homogenate
Assay Type	Quantitative Sandwich
Sensitivity	1.0 ng/mL
Preparation and Storage	Store all reagents at 2-8 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PRL elisa kit	Intended Uses: This PRL ELISA kit is a 1.5 hour solid-phase ELISA designed for the quantitative determination of Porcine PRL. This ELISA kit for research use only, not for therapeutic or test applications! Principle of the Assay: PRL ELISA kit applies the quantitative sandwich enzyme immunoassay technique. The microtiter plate has been pre-coated with a monoclonal antibody specific for PRL. Standards or samples are then added to the microtiter plate wells and PRL if present, will bind to the antibody pre-coated wells. In order to quantitatively determine the amount of PRL present in the sample, a standardized preparation of horseradish peroxidase (HRP)-conjugated polyclonal antibody, specific for PRL are added to each well to "sandwich" the PRL immobilized on the plate. The microtiter plate undergoes incubation, and then the wells are thoroughly washed to remove all unbound components. Next, substrate solutions are added to each well. The enzyme (HRP) and substrate are allowed to react over a short incubation period. Only those wells that contain PRL and enzyme-conjugated antibody will exhibit a change in color. The enzyme-substrate reaction is terminated by addition of a sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450 nm. A standard curve is plotted relating the intensity of the color (O.D.) to the concentration of standards. The PRL concentration in each sample is interpolated from this standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	130930
NCBI GeneID	5617
Molecular Weight	25,876 Da
NCBI Official Full Name	Prolactin
NCBI Official Synonym Full Names	prolactin
NCBI Official Symbol	PRL
NCBI Protein Information	prolactin; decidual prolactin
UniProt Protein Name	Prolactin
UniProt Gene Name	PRL
UniProt Synonym Gene Names	PRL
UniProt Entry Name	PRL_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/prl-porcine-elisa-kits/prolactin/16583>