

Rabbit TBC1D4 Polyclonal Antibody | anti-TBC1D4 antibody

TBC1D4 Polyclonal Antibody

CATALOG # AAA166749	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

TBC1D4; Polyclonal Antibody; TBC1D4 Polyclonal Antibody; AS160; anti-TBC1D4 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	2mg/mL (varies by lot)
Sequence Length	1290
Applicable Applications for anti-TBC1D4 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Synthetic peptide of human TBC1D4
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human brain tissue using TBC1D4 Polyclonal Antibody at dilution 1:30


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TBC1D4 Polyclonal Antibody at dilution 1:30

ADDITIONAL INFORMATION

Related Product Information for anti-TBC1D4 antibody	TBC1 domain family member 4 (TBC1D4), also designated AS160, can be insulin- and/or AKT1-induced. Insulin-stimulated phosphorylation is required for GLUT4 translocation. TBC1D4 may play a role as a GTPase activating protein for proteins in the Rab family. It is expressed primarily in skeletal muscle and heart, as well as spleen, lymph node and leukocytes. Defects in the TBC1D4 gene may cause atopic dermatitis (AD), sometimes referred to as eczema, an atopic chronic skin disease. The skin of affected individuals reacts to irritants or allergens and becomes red, flaky and itchy. The skin is also more vulnerable to inflammations, and symptoms can grow or disappear over time.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	557786188
NCBI GeneID	9882
NCBI Accession #	NP_001273587.1
NCBI GenBank Nucleotide #	NM_001286658.1
Molecular Weight	60,095 Da
NCBI Official Full Name	TBC1 domain family member 4 isoform 2
NCBI Official Synonym Full Names	TBC1 domain family, member 4
NCBI Official Symbol	TBC1D4
NCBI Official Synonym Symbols	AS160; NIDDM5
NCBI Protein Information	TBC1 domain family member 4; Acrg embryonic lethality minimal region ortholog; TBC (Tre-2, BUB2, CDC16) domain-containing protein; akt substrate of 160 kDa
UniProt Protein Name	TBC1 domain family member 4
UniProt Gene Name	TBC1D4
UniProt Synonym Gene Names	AS160; KIAA0603; AS160
UniProt Entry Name	TBCD4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/tbc1d4-antibody-immunohistochemistry-elisa/166749>