

Rabbit anti-Human, Mouse TRAF3 Polyclonal Antibody | anti-TRAF3 antibody

TRAF3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA166919	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TRAF3; Polyclonal Antibody; TRAF3 Antibody; CAP1; LAP1; CAP-1; CRAF1; IIAE5; CD40bp; anti-TRAF3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-TRAF3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human TNF receptor-associated factor 3
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-949 / sc-948 / sc-1828 / sc-1574
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.


IMAGES


IF (Immunofluorescence)

Immunofluorescence analysis of HeLa cell using TRAF3 antibody. Blue: DAPI for nuclear staining.


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human tonsil tissue using TRAF3 antibody.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using TRAF3 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-TRAF3 antibody

The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from, members of the TNF receptor (TNFR) superfamily. This protein participates in the signal transduction of CD40, a TNFR family member important for the activation of the immune response. This protein is found to be a critical component of the lymphotoxin-beta receptor (LTbetaR) signaling complex, which induces NF-kappaB activation and cell death initiated by LTbeta ligation. Epstein-Barr virus encoded latent infection membrane protein-1 (LMP1) can interact with this and several other members of the TRAF family, which may be essential for the oncogenic effects of LMP1. Several alternatively spliced transcript variants encoding three distinct isoforms have been reported.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/traf3-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/166919>