

Rabbit MME Polyclonal Antibody | anti-MME antibody

MME Polyclonal Antibody

CATALOG # AAA167027	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

MME; Polyclonal Antibody; MME Polyclonal Antibody; NEP; SFE; CD10; CALLA; anti-MME antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	1.4mg/mL (varies by lot)
Applicable Applications for anti-MME antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Synthetic peptide of human MME
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.


IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human lung cancer using MME Polyclonal Antibody at dilution of 1:45


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human colon cancer using MME Polyclonal Antibody at dilution of 1:45



WB (Western Blot)

Western Blot analysis of 293T cell using MME Polyclonal Antibody at dilution of 1:700

ADDITIONAL INFORMATION

Related Product Information for anti-MME antibody

This gene encodes a common acute lymphocytic leukemia antigen that is an important cell surface marker in the identification of human acute lymphocytic leukemia (ALL). This protein is present on leukemic cells of pre-B phenotype, which represent 85% of cases of ALL. This protein is not restricted to leukemic cells, however, and is found on a variety of normal tissues. It is a glycoprotein that is particularly abundant in kidney, where it is present on the brush border of proximal tubules and on glomerular epithelium. The protein is a neutral endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon, enkephalins, substance P, neurotensin, oxytocin, and bradykinin. This gene, which encodes a 100-kD type II transmembrane glycoprotein, exists in a single copy of greater than 45 kb. The 5' untranslated region of this gene is alternatively spliced, resulting in four separate mRNA transcripts. The coding region is not affected by alternative splicing.

Product Categories/Family for anti-MME antibody

Cancer; Stem cells

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mme-antibody-immunohistochemistry-western-blot-elisa/167027>