

Rabbit PSMC2 Polyclonal Antibody | anti-PSMC2 antibody

PSMC2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA167079	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

PSMC2; Polyclonal Antibody; PSMC2 Polyclonal Antibody; S7; MSS1; Nbla10058; anti-PSMC2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	2mg/mL (varies by lot)
Sequence Length	433
Applicable Applications for anti-PSMC2 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant protein of human PSMC2
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded Human breast cancer using PSMC2 Polyclonal Antibody at dilution of 1:25



IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded Human thyroid cancer using PSMC2 Polyclonal Antibody at dilution of 1:25



WB (Western Blot)

Western Blot analysis of 231 cell using PSMC2 Polyclonal Antibody at dilution of 1:500

ADDITIONAL INFORMATION

Related Product Information for anti-PSMC2 antibody

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity.

Product Categories/Family for anti-PSMC2 antibody

[Cell Biology](#)

NCBI AND UNIPROT INFORMATION

NCBI GI

[649102561](#)

NCBI Official Full Name

PSMC2, partial

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/psmc2-antibody-immunohistochemistry-western-blot-elisa/167079>