

Rabbit ACY1 Polyclonal Antibody | anti-ACY1 antibody

ACY1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA167576	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

ACY1; Polyclonal Antibody; ACY1 Polyclonal Antibody; ACY-1; ACY1D; anti-ACY1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	0.9mg/mL (varies by lot)
Sequence Length	373
Applicable Applications for anti-ACY1 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant protein of human ACY1
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human colon cancer using ACY1 Polyclonal Antibody at dilution of 1:25



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human gastric cancer using ACY1 Polyclonal Antibody at dilution of 1:25



WB (Western Blot)

Western Blot analysis of Mouse kidney and Human normal kidney tissue, K562 cell using ACY1 Polyclonal Antibody at dilution of 1:250

ADDITIONAL INFORMATION

Related Product Information for anti-ACY1 antibody

This gene encodes a cytosolic, homodimeric, zinc-binding enzyme that catalyzes the hydrolysis of acylated L-amino acids to L-amino acids and an acyl group, and has been postulated to function in the catabolism and salvage of acylated amino acids. This gene is located on chromosome 3p21. 1, a region reduced to homozygosity in small-cell lung cancer (SCLC), and its expression has been reported to be reduced or undetectable in SCLC cell lines and tumors. The amino acid sequence of human aminoacylase-1 is highly homologous to the porcine counterpart, and this enzyme is the first member of a new family of zinc-binding enzymes. Mutations in this gene cause aminoacylase-1 deficiency, a metabolic disorder characterized by central nervous system defects and increased urinary excretion of N-acetylated amino acids. Alternative splicing of this gene results in multiple transcript variants. Read-through transcription also exists between this gene and the upstream ABHD14A (abhydrolase domain containing 14A) gene, as represented in GeneID:100526760. A related pseudogene has been identified on chromosome 18.

NCBI AND UNIPROT INFORMATION

NCBI GI #	312032409
NCBI GeneID	95
NCBI Accession #	NP_001185827.1
NCBI GenBank Nucleotide #	NM_001198898.1
Molecular Weight	42,195 Da
NCBI Official Full Name	aminoacylase-1 isoform d
NCBI Official Synonym Full Names	aminoacylase 1
NCBI Official Symbol	ACY1
NCBI Official Synonym Symbols	ACY-1; ACY1D; HEL-S-5
NCBI Protein Information	aminoacylase-1; N-acyl-L-amino-acid amidohydrolase; epididymis secretory protein Li 5
UniProt Protein Name	Aminoacylase-1
UniProt Gene Name	ACY1
UniProt Synonym Gene Names	ACY-1
UniProt Entry Name	ACY1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/acy1-antibody-immunohistochemistry-western-blot-elisa/167576>