

Rabbit IDH1 Polyclonal Antibody | anti-IDH1 antibody

IDH1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA167945	Rabbit	Human, Mouse, Rat	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

IDH1; Polyclonal Antibody; IDH1 Polyclonal Antibody; IDH; IDP; IDCD; IDPC; PICD; anti-IDH1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	2.2mg/mL (varies by lot)
Sequence Length	414
Applicable Applications for anti-IDH1 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant protein of human IDH1
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.


IMAGES


IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded Human lung cancer using IDH1 Polyclonal Antibody at dilution of 1:50


IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded Human liver cancer using IDH1 Polyclonal Antibody at dilution of 1:50



WB (Western Blot)

Western Blot analysis of 231 and hela cell using IDH1 Polyclonal Antibody at dilution of 1:550

ADDITIONAL INFORMATION

Related Product Information for anti-IDH1 antibody	Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence.
Product Categories/Family for anti-IDH1 antibody	Cancer ; Epigenetics

NCBI AND UNIPROT INFORMATION

NCBI GI #	538917681
NCBI GeneID	3417
NCBI Accession #	NP_001269316.1
NCBI GenBank Nucleotide #	NM_001282387.1
Molecular Weight	46,659 Da
NCBI Official Full Name	isocitrate dehydrogenase
NCBI Official Synonym Full Names	isocitrate dehydrogenase 1 (NADP+), soluble
NCBI Official Symbol	IDH1
NCBI Official Synonym Symbols	IDH; IDP; IDCD; IDPC; PICD; HEL-216; HEL-S-26
NCBI Protein Information	isocitrate dehydrogenase [NADP] cytoplasmic; isocitrate dehydrogenase [NADP] cytoplasmic; NADP(+)-specific ICDH; NADP-dependent isocitrate dehydrogenase, cytosolic; NADP-dependent isocitrate dehydrogenase, peroxisomal; epididymis luminal protein 216; epididymis secretory protein Li 26; oxalosuccinate decarboxylase
UniProt Protein Name	Isocitrate dehydrogenase [NADP] cytoplasmic
UniProt Gene Name	IDH1
UniProt Synonym Gene Names	PICD; IDH
UniProt Entry Name	IDHC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/idh1-antibody-immunohistochemistry-western-blot-elisa/167945>