

Rabbit KLC1 Polyclonal Antibody | anti-KLC1 antibody

KLC1 Polyclonal Antibody

CATALOG # AAA168570	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

KLC1; Polyclonal Antibody; KLC1 Polyclonal Antibody; KLC; KNS2; KNS2A; anti-KLC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	2.2mg/mL (varies by lot)
Sequence Length	618
Applicable Applications for anti-KLC1 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant protein of human KLC1
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human lung cancer using KLC1 Polyclonal Antibody at dilution of 1:40



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human colon cancer using KLC1 Polyclonal Antibody at dilution of 1:40



WB (Western Blot)

Western Blot analysis of Lovo cell using KLC1 Polyclonal Antibody at dilution of 1:550

ADDITIONAL INFORMATION

Related Product Information for anti-KLC1 antibody

Conventional kinesin is a tetrameric molecule composed of two heavy chains and two light chains, and transports various cargos along microtubules toward their plus ends. The heavy chains provide the motor activity, while the light chains bind to various cargos. This gene encodes a member of the kinesin light chain family. It associates with kinesin heavy chain through an N-terminal domain, and six tetratricopeptide repeat (TPR) motifs are thought to be involved in binding of cargos such as vesicles, mitochondria, and the Golgi complex. Thus, kinesin light chains function as adapter molecules and not motors per se. Although previously named "kinesin 2", this gene is not a member of the kinesin-2 / kinesin heavy chain subfamily of kinesin motor proteins. Extensive alternative splicing produces isoforms with different C-termini that are proposed to bind to different cargos; however, the full-length nature and/or biological validity of most of these variants have not been determined.

NCBI AND UNIPROT INFORMATION

NCBI GI #	194306559
NCBI GeneID	3831
NCBI Accession #	NP_001123579.1
NCBI GenBank Nucleotide #	NM_001130107.1
Molecular Weight	69,765 Da
NCBI Official Full Name	kinesin light chain 1 isoform 3
NCBI Official Synonym Full Names	kinesin light chain 1
NCBI Official Symbol	KLC1
NCBI Official Synonym Symbols	KLC; KNS2; KNS2A
NCBI Protein Information	kinesin light chain 1; kinesin 2 60/70kDa; medulloblastoma antigen MU-MB-2.50
UniProt Protein Name	Kinesin light chain 1
UniProt Gene Name	KLC1
UniProt Synonym Gene Names	KLC; KNS2; KLC 1
UniProt Entry Name	KLC1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/klc1-antibody-immunohistochemistry-western-blot-elisa/168570>