Rabbit SETD2 Polyclonal Antibody | anti-SETD2 antibody

SETD2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA168839	**Rabbit**	**Human, Mouse, Rat**	**Immunohistochemistry, ELISA**

PRODUCT SYNONYMS

SETD2; Polyclonal Antibody; SETD2 Polyclonal Antibody; HYPB; SET2; HIF-1; HIP-1; KMT3A; HBP231; HSPC069; p231HBP; anti-SETD2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	1.2mg/mL (varies by lot)
Applicable Applications for anti-SETD2 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant protein of human SETD2
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES



IHC (Immunohiostchemistry)
Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using SETD2 Polyclonal Antibody at dilution 1:30



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human liver cancer tissue using SETD2 Polyclonal Antibody at dilution 1:30

ADDITIONAL INFORMATION

Related Product Information for anti-SETD2 antibody	Huntington's disease (HD), a neurodegenerative disorder characterized by loss of striatal neurons, is caused by an expansion of a polyglutamine tract in the HD protein huntingtin. This gene encodes a protein belonging to a class of huntingtin interacting proteins characterized by WW motifs. This protein is a histone methyltransferase that is specific for lysine-36 of histone H3, and methylation of this residue is associated with active chromatin. This protein also contains a novel transcriptional activation domain and has been found associated with hyperphosphorylated RNA polymerase II.
Product Categories/Family for anti-SETD2 antibody	Cancer; Epigenetics

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: https://stage.aaabiotech.com/polyclonal/setd2-antibody-immunohistochemistry-elisa/168839