

Rabbit SH2D1A Polyclonal Antibody | anti-SH2D1A antibody

SH2D1A Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA168966	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SH2D1A; Polyclonal Antibody; SH2D1A Antibody; DSHP; EBVS; FLJ18687; FLJ92177; IMD5; LYP; MTCP1; SAP; XLP; XLPD; anti-SH2D1A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	1mg/mL (varies by lot)
Sequence Length	125
Applicable Applications for anti-SH2D1A antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human SH2D1A
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-8639 / sc-8638 / sc-8333
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence analysis of HeLa cell using SH2D1A antibody. Blue: DAPI for nuclear staining.




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded rat lung using SH2D1A antibody at dilution of 1:200 (400x lens).



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human stomach cancer using SH2D1A antibody at dilution of 1:200 (400x lens)



WB (Western Blot)
Western blot analysis of extracts of various cell lines, using SH2D1A antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-SH2D1A antibody

SH2D1A and SH2D1B are small, adaptor proteins with a single SH2-domain that play important signal transduction roles mediated by the signaling lymphocytic activation molecule (SLAM) family receptors (1). SH2D1A (also called SAP or SLAM-associated protein) is frequently mutated in patients with X-linked lymphoproliferative disease (Duncan's disease), which is characterized by extreme susceptibility to Epstein-Barr virus; approximately 50 different SH2D1A mutations have been reported to date (2-4). The single SH2D1B gene in humans (also called EAT-2 or Ewing's sarcoma's/FLI1-activated transcript 2) is present as a pair of duplicated EAT-2A and EAT-2B genes with identical genomic organization in mouse and rat (5, 6).

NCBI AND UNIPROT INFORMATION

NCBI GI #	169234945
NCBI GeneID	4068
NCBI Accession #	NP_001108409.1
NCBI GenBank Nucleotide #	NM_001114937.2
Molecular Weight	6,631 Da
NCBI Official Full Name	SH2 domain-containing protein 1A isoform 2
NCBI Official Synonym Full Names	SH2 domain containing 1A
NCBI Official Symbol	SH2D1A
NCBI Official Synonym Symbols	LYP; SAP; XLP; DSHP; EBVS; IMD5; XLPD; MTCP1; XLPD1; SAP/SH2D1A
NCBI Protein Information	SH2 domain-containing protein 1A; Duncan disease SH2-protein; SLAM associated protein/SH2 domain protein 1A; SLAM-associated protein; T cell signal transduction molecule SAP; T-cell signal transduction molecule SAP; signaling lymphocyte activation molecule-associated protein; signaling lymphocytic activation molecule-associated protein
UniProt Protein Name	SH2 domain-containing protein 1A
UniProt Gene Name	SH2D1A
UniProt Synonym Gene Names	DSHP; SAP; SLAM-associated protein
UniProt Entry Name	SH21A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.