

Rabbit ACVR2B Polyclonal Antibody | anti-ACVR2B antibody

ACVR2B Polyclonal Antibody

CATALOG # AAA169715	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

ACVR2B; Polyclonal Antibody; ACVR2B Polyclonal Antibody; HTX4; ACTRIIB; ActR-IIB; anti-ACVR2B antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity purification
Form/Format	PBS with 0.05% sodium azide and 50% glycerol, PH7.4
Applicable Applications for anti-ACVR2B antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant protein of human ACVR2B
Note	Centrifuge before opening to ensure complete recovery of vial contents.
Preparation and Storage	Store at -20°C. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human colon cancer tissue using ACVR2B Polyclonal Antibody at dilution 1:50



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human liver cancer tissue using ACVR2B Polyclonal Antibody at dilution 1:50

ADDITIONAL INFORMATION

Related Product Information for anti-ACVR2B antibody	Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases. This gene encodes activin A type IIB receptor, which displays a 3- to 4-fold higher affinity for the ligand than activin A type II receptor.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	116734708
NCBI GeneID	93
NCBI Accession #	NP_001097.2
NCBI GenBank Nucleotide #	NM_001106.3
UniProt Accession #	Q13705; Q4VAV0
Molecular Weight	58kDa
NCBI Official Full Name	activin receptor type-2B
NCBI Official Synonym Full Names	activin A receptor, type IIB
NCBI Official Symbol	ACVR2B
NCBI Official Synonym Symbols	HTX4; ACTRIIB; ActR-IIB
NCBI Protein Information	activin receptor type-2B
UniProt Protein Name	Activin receptor type-2B
UniProt Gene Name	ACVR2B
UniProt Synonym Gene Names	ACTR-IIB
UniProt Entry Name	AVR2B_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/acvr2b-antibody-immunohistochemistry-elisa/169715>