

Rabbit FABP2 Polyclonal Antibody | anti-FABP2 antibody

FABP2 Polyclonal Antibody

CATALOG # AAA170185	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

FABP2; Polyclonal Antibody; FABP2 Polyclonal Antibody; FABP1; I-FABP; anti-FABP2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	2.5mg/mL (varies by lot)
Sequence Length	132
Applicable Applications for anti-FABP2 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Synthetic peptide of human FABP2
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human brain using FABP2 Polyclonal Antibody at dilution of 1:40



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human liver cancer using FABP2 Polyclonal Antibody at dilution of 1:40



WB (Western Blot)

Western Blot analysis of Mouse small intestine tissue using FABP2 Polyclonal Antibody at dilution of 1:650

ADDITIONAL INFORMATION

Related Product Information for anti-FABP2 antibody

The intracellular fatty acid-binding proteins (FABPs) belong to a multigene family with nearly twenty identified members. FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa proteins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fatty acids. They may also be responsible in the modulation of cell growth and proliferation. Intestinal fatty acid-binding protein 2 gene contains four exons and is an abundant cytosolic protein in small intestine epithelial cells. This gene has a polymorphism at codon 54 that identified an alanine-encoding allele and a threonine-encoding allele. Thr-54 protein is associated with increased fat oxidation and insulin resistance.

Product Categories/Family for anti-FABP2 antibody

Stem Cells

NCBI AND UNIPROT INFORMATION

NCBI GI #	194097325
NCBI GeneID	2169
NCBI Accession #	NP_000125.2
NCBI GenBank Nucleotide #	NM_000134.3
Molecular Weight	15,207 Da
NCBI Official Full Name	fatty acid-binding protein, intestinal
NCBI Official Synonym Full Names	fatty acid binding protein 2, intestinal
NCBI Official Symbol	FABP2
NCBI Official Synonym Symbols	FABPI; I-FABP
NCBI Protein Information	fatty acid-binding protein, intestinal; fatty acid-binding protein 2; intestinal-type fatty acid-binding protein
UniProt Protein Name	Fatty acid-binding protein, intestinal
UniProt Gene Name	FABP2
UniProt Synonym Gene Names	FABPI; I-FABP
UniProt Entry Name	FABPI_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/fabp2-antibody-immunohistochemistry-western-blot-elisa/170185>