

Rabbit SEPT7 Polyclonal Antibody | anti-SEPT7 antibody

SEPT7 Polyclonal Antibody

CATALOG # AAA170376	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	--	--


PRODUCT SYNONYMS

SEPT7; Polyclonal Antibody; SEPT7 Polyclonal Antibody; CDC3; CDC10; SEPT7A; NBLA02942; anti-SEPT7 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen affinity purification
Concentration	2.3mg/mL (varies by lot)
Applicable Applications for anti-SEPT7 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant protein of human SEPT7
Buffer	PBS with 0.05% sodium azide, 50% glycerol, pH7.3
Preparation and Storage	Store at -20 degree C (regular) and -80 degree C (long term). Avoid freeze / thaw cycles.

IMAGES




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded Human gastric cancer using SEPT7 Polyclonal Antibody at dilution of 1:50



IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded Human colon cancer using SEPT7 Polyclonal Antibody at dilution of 1:50



WB (Western Blot)

Western Blot analysis of Hela cell using SEPT7 Polyclonal Antibody at dilution of 1:380

ADDITIONAL INFORMATION

Related Product Information for anti-SEPT7 antibody	This gene encodes a protein that is highly similar to the CDC10 protein of Saccharomyces cerevisiae. The protein also shares similarity with Diff 6 of Drosophila and with H5 of mouse. Each of these similar proteins, including the yeast CDC10, contains a GTP-binding motif. The yeast CDC10 protein is a structural component of the 10 nm filament which lies inside the cytoplasmic membrane and is essential for cytokinesis. This human protein functions in gliomagenesis and in the suppression of glioma cell growth, and it is required for the association of centromere-associated protein E with the kinetochore. Alternative splicing results in multiple transcript variants. Several related pseudogenes have been identified on chromosomes 5, 7, 9, 10, 11, 14, 17 and 19
---	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/sept7-antibody-immunohistochemistry-western-blot-elisa/170376>