

Monkey Soluble transferrin receptor ELISA Kit | sTfR elisa kit

Monkey Soluble transferrin receptor ELISA Kit

CATALOG #
AAA17165

REACTIVITY
Monkey


PRODUCT SYNONYMS

Soluble transferrin receptor; Monkey Soluble transferrin receptor ELISA Kit; sTfR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Monkey
Specificity	This assay has high sensitivity and excellent specificity for detection of sTFER. No significant cross-reactivity or interference between sTFER and analogues was observed. NOTE: Limited by current skills and knowledge, it is impossible for us to complete the cross-reactivity detection between sTFER and all the analogues, therefore, cross reaction may still exist in some cases.
Assay Type	Quantitative Sandwich
Samples	Serum, plasma, cell culture supernatants, body fluid and tissue homogenate
Sensitivity	1.0 ng/mL
Preparation and Storage	Store all reagents at 2-8 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for sTfR elisa kit	Principle of the Assay: sTfEr ELISA kit applies the quantitative sandwich enzyme immunoassay technique. The microtiter plate has been pre-coated with a monoclonal antibody specific for sTfEr. Standards or samples are then added to the microtiter plate wells and sTfEr if present, will bind to the antibody pre-coated wells. In order to quantitatively determine the amount of sTfEr present in the sample, a standardized preparation of horseradish peroxidase (HRP)-conjugated polyclonal antibody, specific for sTfEr are added to each well to "sandwich" the sTfEr immobilized on the plate. The microtiter plate undergoes incubation, and then the wells are thoroughly washed to remove all unbound components. Next, substrate solutions are added to each well. The enzyme (HRP) and substrate are allowed to react over a short incubation period. Only those wells that contain sTfEr and enzyme-conjugated antibody will exhibit a change in color. The enzyme-substrate reaction is terminated by addition of a sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450 nm. A standard curve is plotted relating the intensity of the color (O.D.) to the concentration of standards. The sTfEr concentration in each sample is interpolated from this standard curve. Intended Uses: This sTfEr ELISA kit is a 1.5 hour solid-phase ELISA designed for the quantitative determination of Monkey sTfEr. This ELISA kit for research use only, not for therapeutic or diagnostic applications!
Product Categories/Family for sTfR elisa kit	Signal Transduction

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/stfr-monkey-elisa-kits/soluble-transferrin-receptor/17165>