

ALK-2 / ACVR1 / ALK2 recombinant protein

Recombinant Human ALK-2 / ACVR1 / ALK2 Protein (His tag)

CATALOG # AAA173509	HOST Baculovirus-Insect Cells	REACTIVITY Human
-------------------------------	---	----------------------------


PRODUCT SYNONYMS

ALK-2 / ACVR1 / ALK2; Recombinant Human ALK-2 / ACVR1 / ALK2 Protein (His tag); Activin Receptor Type-1; Activin Receptor Type I; ACTR-I; Activin Receptor-Like Kinase 2; ALK-2; Serine/Threonine-Protein Kinase Receptor R1; SKR1; TGF-B Superfamily Receptor Type I; TSR-I; ACVR1; ACVRLK2;ACVR1A;ACVRLK2;ALK2;FOP;SKR1; ALK-2 / ACVR1 / ALK2 recombinant protein

PRODUCT SPECIFICATIONS

Host	Baculovirus-Insect Cells
Reactivity	Human
Purity/Purification	>93 % as determined by reducing SDS-PAGE.
Form/Format	Lyophilized from sterile 20mM Tris, 500mM NaCl, 25% glycerol, 5% Trehalose, 5% Mannitol, 0.01 % Tween-80, pH 7.5.
Sequence	Met1-Val12 4
Predicted N Terminal	Met 21
Species	Human
Tag	C-His
Endotoxin	< 1.0 EU per ug as determined by the LAL method.
Reconstitution	We recommend that the vial be centrifuged to opening to bring the contents to the bottom. Do not mix by vortex. It is recommended that sterile water be added to the vial to prepare a stock solution of 0.25mg/mL. Concentration is measured by UV-Vis.
Preparation and Storage	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for ALK-2 / ACVR1 / ALK2 recombinant protein	Background: ALK-2, also termed as ACVR1, was initially identified as an activin type I receptor because of its ability to bind activinin concert with ActRII or ActRIIB. ALK-2 is also identified as a BMP type I receptor. It has been demonstrated that ALK-2 forms complex with either the BMP-2/7-bound BMPR-II or ACVR2A /ACVR2B. ALK-1 and ALK-2 presentingin the yeast Saccharomyces cerevisiae are two haspin homologues. Both ALK-1 and ALK-2 exhibit a weak auto-kinase activity in vitro, and are phosphoproteins in vivo. ALK-1 and ALK-2 levels peak in mitosis and late-S/G2. Control of protein stability plays a major role in ALK-2 regulation. The half-life of ALK-2 is particularly short in G1. Over expression of ALK-2, but not of ALK-1, causes a mitotic arrest, which is correlated to the kinase activity of the protein. This suggests a role for ALK-2 in the control of mitosis. Endoglin is phosphorylated on cytosolic domainthreonine residues by the TGF-beta type I receptors ALK-2 and ALK-5 in prostate cancer cells. Endoglin did not inhibit cell migration in the presence of constitutively active ALK-2. Defects in ALK-2 are a cause of fibrodysplasia ossificans progressiva (FOP).
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4501895
NCBI GeneID	90
NCBI Accession #	NP_001096.1
NCBI GenBank Nucleotide #	NM_001105.4
UniProt Accession #	Q04771
Molecular Weight	Mol_Mass: 12.8 kDa AP_Mol_Mass: 17 kDa
NCBI Official Full Name	activin receptor type-1
NCBI Official Synonym Full Names	activin A receptor type 1
NCBI Official Symbol	ACVR1
NCBI Official Synonym Symbols	FOP; ALK2; SKR1; TSRI; ACTRI; ACVR1A; ACVRLK2
NCBI Protein Information	activin receptor type-1
UniProt Protein Name	Activin receptor type-1
UniProt Gene Name	ACVR1
UniProt Synonym Gene Names	ACVRLK2; ACTR-I; ALK-2; SKR1; TSR-I

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/acvr1-recombinant-protein-baculovirus-insect-cells/173509>