

GNMB / Osteoactivin Recombinant Protein | GNMB recombinant protein

Recombinant Human GNMB Protein (aa 1-474, His Tag)

CATALOG # AAA173548	HOST HEK293 Cells
-------------------------------	-----------------------------


PRODUCT SYNONYMS

GNMB / Osteoactivin; Recombinant Human GNMB Protein (aa 1-474; His Tag); Transmembrane Glycoprotein NMB; Transmembrane Glycoprotein HGFIN; GNMB; HGFIN; NMB; Osteoactivin; GNMB recombinant protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	> 98 % as determined by reducing SDS-PAGE.
Form/Format	Lyophilized from sterile PBS, pH 7.4, 5% Trehalose, 5% Mannitol, 0.01% Tween-80.
Sequence	Met1-Pro47 4
Endotoxin	< 1.0 EU per ug of the protein as determined by the LAL method.
Species	Human
Tag	C-His
Reconstitution	We recommend that the vial be centrifuged to opening to bring the contents to the bottom. Do not mix by vortex. It is recommended that sterile water(200ul) be added to the vial to prepare a stock solution of 0.25mg/mL. Concentration is measured by UV-Vis.
Preparation and Storage	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80 degree C . Reconstituted protein solution can be stored at 4-8 degree C for 2-7 days. Aliquots of reconstituted samples are stable at < -20 degree C for 3 months. Shipping: This product is provided as lyophilized powder which is shipped with blue ice.

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for GPNMB recombinant protein	Background: GPNMB belongs to the PMEL / NMB family, also known as Osteoactivin and Hematopoietic growth factor-inducible neurokinin 1 (HGFIN), is a transmembrane glycoprotein that is expressed in numerous cells, including osteoclasts, macrophages, dendritic cells, and tumor cells. It is suggested to influence osteoblast maturation, cell adhesion and migration. GPNMB protein acts as a downstream mediator of BMP-2 effects on osteoblast differentiation and function. GPNMB participates in bone mineralization, and functions as a negative regulator of inflammation in macrophages. Osteoactivin is expressed at high levels in normal and inflammatory liver macrophages suggesting a significant role in acute liver injury. The early-phase upregulation of Osteoactivin expression in the tubular epithelium in response to renal injury might play a role in triggering renal interstitial fibrosis via activation of matrix metalloproteinase expression and collagen remodeling in rats. Osteoactivin as a protein that is expressed in aggressive human breast cancers and is capable of promoting breast cancer metastasis to bone.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	52694752
NCBI GeneID	10457
NCBI Accession #	NP_001005340.1
NCBI GenBank Nucleotide #	NM_001005340.1
UniProt Accession #	Q14956; Q6UVX1; Q8N1A1; A4D155
Molecular Weight	52.1 kDa
NCBI Official Full Name	transmembrane glycoprotein NMB isoform a
NCBI Official Synonym Full Names	glycoprotein nmb
NCBI Official Symbol	GPNMB
NCBI Official Synonym Symbols	NMB; HGFIN
NCBI Protein Information	transmembrane glycoprotein NMB
UniProt Protein Name	Transmembrane glycoprotein NMB
UniProt Gene Name	GPNMB
UniProt Synonym Gene Names	HGFIN; NMB

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/gpnmb-recombinant-protein-hek293-cells/173548>