

Rabbit anti-Human MEK2 Polyclonal Antibody | anti-MEK2 antibody

MEK2 Polyclonal Antibody

CATALOG # AAA173860	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

MEK2; Polyclonal Antibody; MEK2 Polyclonal Antibody; Dual specificity mitogen-activated protein kinase kinase 2 (MAP kinase kinase 2) (MAPKK 2) (EC 2.7.12.2) (ERK activator kinase 2) (MAPK/ERK kinase 2) (MEK 2); anti-MEK2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	1mg/mL (varies by lot)
Sequence Length	400
Applicable Applications for anti-MEK2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant Protein of MEK2
Buffer	PBS with 0.02% sodium azide and 50% glycerol pH 7.4.
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human colon carcinoma tissue using MEK2 Polyclonal Antibody at dilution of 1:200.



WB (Western Blot)

Western Blot analysis of 1) Hela, 2) 293T cells using MEK2 Polyclonal Antibody at dilution of 1:2000.

ADDITIONAL INFORMATION

Related Product Information for anti-MEK2 antibody	The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	13489054
NCBI GeneID	5605
NCBI Accession #	NP_109587.1
NCBI GenBank Nucleotide #	NM_030662.3
UniProt Accession #	P36507
Molecular Weight	45kDa
NCBI Official Full Name	dual specificity mitogen-activated protein kinase kinase 2
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 2
NCBI Official Symbol	MAP2K2
NCBI Official Synonym Symbols	CFC4; MEK2; MKK2; MAPKK2; PRKMK2
NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 2
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 2
UniProt Gene Name	MAP2K2
UniProt Synonym Gene Names	MEK2; MKK2; PRKMK2; MAP kinase kinase 2; MAPKK 2; MEK 2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mek2-antibody-immunohistochemistry-western-blot-elisa/173860>