

CD44/MIC4 recombinant protein

Recombinant Human CD44/MIC4 Protein (C-6His)

CATALOG # AAA174009	HOST Human Cells
-------------------------------	----------------------------

PRODUCT SYNONYMS

CD44/MIC4; Recombinant Human CD44/MIC4 Protein (C-6His); CD44 Antigen; CDw44; Epican; Extracellular Matrix Receptor III; ECMR-III; GP90 Lymphocyte Homing/Adhesion Receptor; HUTCH-I; Heparan Sulfate Proteoglycan; Hermes Antigen; Hyaluronate Receptor; Phagocytic Glycoprotein 1; PGP-1; Phagocytic Glycoprotein I; PGP-I; CD44; LHR; MDU2; MDU3; MIC4; CD44/MIC4 recombinant protein

PRODUCT SPECIFICATIONS

Host	Human Cells
Purity/Purification	>95% as determined by reducing SDS-PAGE.
Form/Format	Lyophilized from a 0.2 ?m filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Sequence Positions	Gln21-Pro220
Sequence Length	742
Species	Human
Endotoxin	<1.0 EU per ug as determined by LAL test.
Tag	C-6His
Reconstitution	We recommend that the vial be centrifuged to prior to opening to bring contents to the bottom. Do not mix by vortex. It is recommended that the sterile water be added to the vial to prepare a stock solution of 0.25 mg/mL. Concentration is measured by UV-Vis
Preparation and Storage	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

ADDITIONAL INFORMATION

Related Product Information for CD44/MIC4 recombinant protein	CD44 is a cell-surface receptor for hyaluronic acid and also interacts with other ligands, such as osteopontin, collagens, and matrix metalloproteinases. A large number of CD44 isoforms can be generated by the insertion of different combinations of at least nine exons. Increased CD44 antigen is associated with relapses in non-small cell lung cancers. Furthermore, an increasing quantity of evidence suggests that CD44 has various functions related to inflammatory disease. CD44 deficiency induces severe liver injury. CD44-hyaluronate mediates in lymphocyte T and monocyte adhesion to the endothelium, stimulates proinflammatory cytokine release from macrophages and participates in dedifferentiation phenotype of smooth muscle cells from contractile state to synthetic one.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	48255935
NCBI GeneID	960
NCBI Accession #	NP_000601.3
NCBI GenBank Nucleotide #	NM_000610.3
UniProt Accession #	P16070; O95370; P22511; Q04858; Q13419; Q13957; Q13958; Q13959; A5YRN9; B6EAT9; D3DR12; D3DR13

Molecular Weight	Molecular Mass: 23.1 kDa Actual Protein Molecular Mass: 38-50 kDa
NCBI Official Full Name	CD44 antigen isoform 1
NCBI Official Synonym Full Names	CD44 molecule (Indian blood group)
NCBI Official Symbol	CD44
NCBI Official Synonym Symbols	IN; LHR; MC56; MDU2; MDU3; MIC4; Pgp1; CDW44; CSPG8; HCELL; HUTCH-I; ECMR-III
NCBI Protein Information	CD44 antigen
UniProt Protein Name	CD44 antigen
UniProt Gene Name	CD44
UniProt Synonym Gene Names	LHR; MDU2; MDU3; MIC4; ECMR-III; PGP-1; PGP-I

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/cd44-recombinant-protein-human-cells/174009>