

Rabbit anti-Human S100A8 Polyclonal Antibody | anti-S100A8 antibody

S100A8 Polyclonal Antibody

CATALOG # AAA174077	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

S100A8; Polyclonal Antibody; S100A8 Polyclonal Antibody; 60B8Ag; AI323541; B8Ag; BEE11; CAGA; Calgranulin-A; Calprotectin L1L subunit; Calprotectin; included; CFAG; CGLA; Chemotactic cytokine CP-10; CP-10; Cystic fibrosis antigen; L1Ag; Leukocyte L1 complex light chain; MA387; MIF; Migration inhibitory factor-related protein 8; MRP-8; Myeloid-related protein 8; Neutrophil cytosolic 7 kDa protein; NIF; p8; Pro-inflammatory S100 cytokine; Protein S100-A8; S100 calcium binding protein A8 (calgranulin A); S100 calcium binding protein A8; S100 calcium-binding protein A8; S100A8/S100A9 complex; S10A8; Urinary stone protein band A; anti-S100A8 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Concentration	0.36 mg/mL (varies by lot)
Applicable Applications for anti-S100A8 antibody	IHC (Immunohistochemistry)
Immunogen	Recombinant Human Protein S100-A8 protein
Conjugation	Unconjugated
Buffer	PBS with 0.05% Proclin300 and 50% glycerol, pH7.4.
Preparation and Storage	Store at -20 degree C. Avoid freeze / thaw cycles.

IMAGES



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human cervix using S100A8 Polyclonal Antibody at dilution of 1:100(400x)



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded Human cervix using S100A8 Polyclonal Antibody at dilution of 1:100(100x)

ADDITIONAL INFORMATION

Related Product Information for anti-S100A8 antibody	The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in the inhibition of casein kinase and as a cytokine. Altered expression of this protein is associated with the disease cystic fibrosis. Multiple transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-S100A8 antibody	Cancer; Neuroscience

NCBI AND UNIPROT INFORMATION

UniProt Accession #	P05109
---------------------	--------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/s100a8-antibody-immunohistochemistry-elisa/174077>