

Rabbit anti-Human, Mouse COL1A1 Polyclonal Antibody | anti-COL1A1 antibody

COL1A1 Polyclonal Antibody

CATALOG # AAA174138	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	-----------------------------------	---


PRODUCT SYNONYMS

COL1A1; Polyclonal Antibody; COL1A1 Polyclonal Antibody; Alpha 1 type I collagen; Alpha-1 type I collagen; CO1A1; Collagen alpha-1(I) chain; Collagen type I alpha 1; Procollagen type I C terminal propeptide; anti-COL1A1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Concentration	1 mg/mL (varies by lot)
Sequence Length	1,464
Applicable Applications for anti-COL1A1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide of human COL1A1
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded human esophagus using COL1A1 antibody.



IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded human prostate cancer using COL1A1 antibody.



WB (Western Blot)

Western blot analysis of extracts of mouse heart, using COL1A1 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-COL1A1 antibody

This gene encodes the pro-alpha1 chains of type I collagen whose triple helix comprises two alpha1 chains and one alpha2 chain. Type I is a fibril-forming collagen found in most connective tissues and is abundant in bone, cornea, dermis and tendon. Mutations in this gene are associated with osteogenesis imperfecta types I-IV, Ehlers-Danlos syndrome type VIIA, Ehlers-Danlos syndrome Classical type, Caffey Disease and idiopathic osteoporosis. Reciprocal translocations between chromosomes 17 and 22, where this gene and the gene for platelet-derived growth factor beta are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans, resulting from unregulated expression of the growth factor. Two transcripts, resulting from the use of alternate polyadenylation signals, have been identified for this gene.

NCBI AND UNIPROT INFORMATION

NCBI GI #	110349772
NCBI GeneID	1277
NCBI Accession #	NP_000079.2
NCBI GenBank Nucleotide #	NM_000088.3
UniProt Accession #	P02452; O76045; P78441; Q13896; Q13902; Q13903; Q14037; Q14992; Q15176; Q15201; Q16050; Q59F64
Molecular Weight	Calculated: 138kDa Observed: 95kDa; 110kDa
NCBI Official Full Name	collagen alpha-1(I) chain preproprotein
NCBI Official Synonym Full Names	collagen type I alpha 1 chain
NCBI Official Symbol	COL1A1
NCBI Official Synonym Symbols	OI1; OI2; OI3; OI4; EDSC; EDSARTH1
NCBI Protein Information	collagen alpha-1(I) chain
UniProt Protein Name	Collagen alpha-1(I) chain
UniProt Gene Name	COL1A1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/col1a1-antibody-immunohistochemistry-western-blot-elisa/174138>