

Rabbit anti-Human, Mouse CST3 Polyclonal Antibody | anti-CST3 antibody

CST3 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA174204	Rabbit	Human, Mouse	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CST3; Polyclonal Antibody; CST3 Polyclonal Antibody; AD 8; AD8; Amyloid angiopathy and cerebral hemorrhage; ARMD11; bA218C14.4 (cystatin C); bA218C14.4; Cst 3; Cst3; CST3 protein; Cystatin 3; Cystatin-3; Cystatin-C; Cystatin3; CystatinC; CYTC; Epididymis secretory protein Li 2; Gamma trace; Gamma-trace; HCCAA; HEL S 2; MGC117328; Neuroendocrine basic polypeptide; Post gamma globulin; Post-gamma-globulin; anti-CST3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purification
Form/Format	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Concentration	1 mg/mL (varies by lot)
Sequence Length	146
Applicable Applications for anti-CST3 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant protein of human CST3
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human prostate using CST3 antibody.



WB (Western Blot)

Western blot analysis of extracts of various cell lines, using CST3 antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-CST3 antibody	The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes the most abundant extracellular inhibitor of cysteine proteases, which is found in high concentrations in biological fluids and is expressed in virtually all organs of the body. A mutation in this gene has been associated with amyloid angiopathy. Expression of this protein in vascular wall smooth muscle cells is severely reduced in both atherosclerotic and aneurysmal aortic lesions, establishing its role in vascular disease. In addition, this protein has been shown to have an antimicrobial function, inhibiting the replication of herpes simplex virus. Alternative splicing results in multiple transcript variants encoding a single protein.
Product Categories/Family for anti-CST3 antibody	Cancer; Cell Biology; Cardiovascular; Stem cells; Tags & Cell Markers

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503107
NCBI GeneID	1471
NCBI Accession #	NP_000090.1
NCBI GenBank Nucleotide #	NM_000099.4
UniProt Accession #	P01034; Q6FGW9; B2R5J9; D3DW42
Molecular Weight	Calculated: 15kDa Observed: 15kDa
NCBI Official Full Name	cystatin-C
NCBI Official Synonym Full Names	cystatin C
NCBI Official Symbol	CST3
NCBI Official Synonym Symbols	ARMD11; HEL-S-2
NCBI Protein Information	cystatin-C
UniProt Protein Name	Cystatin-C
UniProt Gene Name	CST3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cst3-antibody-immunohistochemistry-western-blot-elisa/174204>