

Rabbit anti-Human, Mouse USO1 Polyclonal Antibody | anti-USO1 antibody

USO1 Polyclonal Antibody

CATALOG # AAA175206	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS ELISA, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	-----------------------------------	--


PRODUCT SYNONYMS

USO1; Polyclonal Antibody; USO1 Polyclonal Antibody; General vesicular transport factor; General vesicular transport factor p115; P115; Protein USO1 homolog; TAP; Transcytosis associated protein; Transcytosis-associated protein; Uso1; VDP; Vesicle docking protein; Vesicle-docking protein; anti-USO1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purification
Form/Format	PBS with 0.05% NaN3 and 40% Glycerol, pH7.4
Concentration	0.4mg/mL (varies by lot)
Sequence Length	962
Applicable Applications for anti-USO1 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Fusion protein of human USO1
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES




IHC (Immunohistochemisry)

Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using USO1 Polyclonal Antibody at dilution of 1:30(×200



IHC (Immunohiostchemistry)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using USO1 Polyclonal Antibody at dilution of 1:30(×200



WB (Western Blot)

Western blot analysis of Mouse testis tissue Jurkat and A549 cell lysates using USO1 Polyclonal Antibody at dilution of 1:350

ADDITIONAL INFORMATION

Related Product Information for anti-USO1 antibody	The protein encoded by this gene is a peripheral membrane protein which recycles between the cytosol and the Golgi apparatus during interphase. It is regulated by phosphorylation: dephosphorylated protein associates with the Golgi membrane and dissociates from the membrane upon phosphorylation. Ras-associated protein 1 recruits this protein to coat protein complex II (COPII) vesicles during budding from the endoplasmic reticulum, where it interacts with a set of COPII vesicle-associated SNAREs to form a cis-SNARE complex that promotes targeting to the Golgi apparatus. Alternative splicing results in multiple transcript variants.
Product Categories/Family for anti-USO1 antibody	Cancer; Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	143743108
NCBI GeneID	8615
UniProt Accession #	O60763
Molecular Weight	108 kDa
NCBI Official Full Name	General vesicular transport factor p115
NCBI Official Synonym Full Names	USO1 vesicle transport factor
NCBI Official Symbol	USO1
NCBI Official Synonym Symbols	TAP; VDP; P115
NCBI Protein Information	general vesicular transport factor p115
UniProt Protein Name	General vesicular transport factor p115
UniProt Gene Name	USO1
UniProt Synonym Gene Names	VDP; TAP
UniProt Entry Name	USO1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/uso1-antibody-immunohistochemistry-western-blot-elisa/175206>