

Rabbit anti-Human IGHA1 Polyclonal Antibody | anti-IGHA1 antibody

IGHA1 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA175338	Rabbit	Human	ELISA, Immunohistochemistry


PRODUCT SYNONYMS

IGHA1; Polyclonal Antibody; IGHA1 Polyclonal Antibody; FLJ14473; FLJ35065; FLJ35500; FLJ36402; FLJ39698; FLJ40001; FLJ41548; FLJ41552; FLJ41789; FLJ43248; FLJ43594; FLJ44293; FLJ46028; FLJ46621; FLJ46724; FLJ46811; FLJ46824; FLJ90170; Ig alpha 1 chain C region; IgA1; Immunoglobulin heavy constant alpha 1; MGC102857; anti-IGHA1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purification
Form/Format	PBS with 0.05% NaN ₃ and 40% Glycerol, pH7.4
Concentration	1.14mg/mL (varies by lot)
Sequence Length	353
Applicable Applications for anti-IGHA1 antibody	ELISA, IHC (Immunohistochemistry)
Immunogen	Fusion protein of human IGHA1
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using IGHA1 Polyclonal Antibody at dilution of 1:65(×200)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using IGHA1 Polyclonal Antibody at dilution of 1:65(×200)

ADDITIONAL INFORMATION

Related Product Information for anti-IGHA1 antibody	Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:22158414, PubMed:20176268). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268). Ig alpha is the major immunoglobulin class in body secretions (PubMed:2241915).
Product Categories/Family for anti-IGHA1 antibody	Cancer; Immunology

NCBI AND UNIPROT INFORMATION

NCBI GI #	113584
NCBI GeneID	3493
UniProt Accession #	P01876
NCBI Official Full Name	Immunoglobulin heavy constant alpha 1
NCBI Official Synonym Full Names	immunoglobulin heavy constant alpha 1
NCBI Official Symbol	IGHA1
NCBI Official Synonym Symbols	IgA1
UniProt Protein Name	Ig alpha-1 chain C region
UniProt Gene Name	IGHA1
UniProt Entry Name	IGHA1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.