

Rabbit SCP2 Polyclonal Antibody | anti-SCP2 antibody

SCP2 Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA176147	Rabbit	Human, Mouse, Rat	ELISA, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SCP2; Polyclonal Antibody; SCP2 Polyclonal Antibody; DKFZp686C12188; DKFZp686D11188; NLTP; Non-specific lipid-transfer protein; Nonspecific lipid transfer protein; NSL TP; NSL-TP; OTTHUMP00000010488; OTTHUMP00000231766; OTTHUMP00000231767; OTTHUMP00000231768; OTTHUMP00000231769; OTTHUMP00000231770; OTTHUMP00000231772; OTTHUMP00000231773; OTTHUMP00000231774; OTTHUMP00000231776; OTTHUMP00000234662; Propanoyl CoA C acyltransferase; Propanoyl-CoA C-acyltransferase; SCP 2; SCP chi; SCP X; SCP-2; SCP-chi; SCP-X; SCPchi; SCPX; Sterol carrier protein 2; Sterol carrier protein X; anti-SCP2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purification
Form/Format	PBS with 0.05% NaN ₃ and 40% Glycerol, pH7.4
Concentration	0.6mg/mL (varies by lot)
Sequence Length	143
Applicable Applications for anti-SCP2 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Fusion protein of human SCP2
Conjugation	Unconjugated
Preparation and Storage	Store at -20 degree C. Avoid freeze/thaw cycles.

IMAGES


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded Human liver cancer tissue using SCP2 Polyclonal Antibody at dilution of 1:35(×200)


WB (Western Blot)

Western blot analysis of Mouse testis tissue lysate using SCP2 Polyclonal Antibody at dilution of 1:350

ADDITIONAL INFORMATION

Related Product Information for anti-SCP2 antibody	This gene encodes two proteins: sterol carrier protein X (SCPx) and sterol carrier protein 2 (SCP2), as a result of transcription initiation from 2 independently regulated promoters. The transcript initiated from the proximal promoter encodes the longer SCPx protein, and the transcript initiated from the distal promoter encodes the shorter SCP2 protein, with the 2 proteins sharing a common C-terminus. Evidence suggests that the SCPx protein is a peroxisome-associated thiolase that is involved in the oxidation of branched chain fatty acids, while the SCP2 protein is thought to be an intracellular lipid transfer protein. This gene is highly expressed in organs involved in lipid metabolism, and may play a role in Zellweger syndrome, in which cells are deficient in peroxisomes and have impaired bile acid synthesis. Alternative splicing of this gene produces multiple transcript variants, some encoding different isoforms.
Product Categories/Family for anti-SCP2 antibody	Cancer; Metabolism; Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	263551
NCBI GeneID	6342
UniProt Accession #	P22307
Molecular Weight	59 kDa
NCBI Official Full Name	sterol carrier protein 2
NCBI Official Synonym Full Names	sterol carrier protein 2
NCBI Official Symbol	SCP2
NCBI Official Synonym Symbols	NLTP; SCPX; SCP-2; SCP-X; NSL-TP; SCP-CHI
NCBI Protein Information	non-specific lipid-transfer protein
UniProt Protein Name	Non-specific lipid-transfer protein
UniProt Gene Name	SCP2
UniProt Synonym Gene Names	NSL-TP; SCP-2; SCP-X
UniProt Entry Name	NLTP_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/scp2-antibody-immunohistochemistry-western-blot-elisa/176147>