

Tie-2 recombinant protein

Recombinant Human Tie-2 (C-6His)

CATALOG #
AAA177961

HOST
Human Cells


PRODUCT SYNONYMS

Tie-2; Recombinant Human Tie-2 (C-6His); angiopoietin-1 receptor; CD202b antigen; CD202b; EC 2.7.10; EC 2.7.10.1; hTIE2; p140 TEK; soluble TIE2 variant 1; soluble TIE2 variant 2; TEK tyrosine kinase; endothelial; TEK; Tie2; VMCM; VMCM1; Tie-2 recombinant protein

PRODUCT SPECIFICATIONS

Host	Human Cells
Purity/Purification	>95% as determined by reducing SDS-PAGE
Form/Format	Lyophilized from a 0.2um filtered solution of PBS, pH7.4.
Sequence	Ala23-Lys7 45
Species	Human
Tag	C-6His
Endotoxin	<1.0EU per ug as determined by the LAL method.
Reconstitution	We recommend that the vial be centrifuged to opening to bring the contents to the bottom. Do not mix by vortex. It is recommended that sterile water be added to the vial to prepare a stock solution of 0.25 mg/mL. Concentration is measured by UV-Vis.
Preparation and Storage	Lyophilized protein should be stored at < -20°C. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable < -20°C for 3 months

IMAGES



SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for Tie-2 recombinant protein	Tie-1/Tie (tyrosine kinase with Ig and EGF homology domains 1) and Tie-2/Tek comprise a receptor tyrosine kinase (RTK) subfamily with unique structural characteristics: two immunoglobulin-like domains flanking three epidermal growth factor (EGF)-like domains and followed by three fibronectin type III-like repeats in the extracellular region and a split tyrosine kinase domain in the cytoplasmic region. These receptors are expressed primarily on endothelial and hematopoietic progenitor cells and play critical roles in angiogenesis, vasculogenesis and hematopoiesis. Human Tie-2 cDNA encodes a 1124 amino acid (aa) residue precursor protein with an 18 residue putative signal peptide, a 727 residue extracellular domain and a 354 residue cytoplasmic domain. Two ligands, angiopoietin-1 (Ang-1) and angiopoietin-2 (Ang-2), which bind Tie-2 with high-affinity have been identified. Ang-2 has been reported to act as an antagonist for Ang-1. Mice engineered to overexpress Ang-2 or to lack Ang-1 or Tie-2 display similar angiogenesis defects.
Product Categories/Family for Tie-2 recombinant protein	Signal Transduction; Cardiovascular; Cancer; Developmental Biology; Stem cells

NCBI AND UNIPROT INFORMATION

NCBI GI #	88758596
NCBI GeneID	7010
NCBI Accession #	NP_000450.2
NCBI GenBank Nucleotide #	NM_000459.3
UniProt Accession #	Q02763; Q5TCU2; Q8IV34; A8K6W0; B4DH20; B4DHD3; D3DRK5; E7EWI2
Molecular Weight	Molecular Mass: 81.4 kDa Actual Protein Molecular Mass: 100-120 kDa
NCBI Official Full Name	angiopoietin-1 receptor
NCBI Official Synonym Full Names	TEK tyrosine kinase, endothelial
NCBI Official Symbol	TEK
NCBI Official Synonym Symbols	TIE2; VMCM; TIE-2; VMCM1; CD202B
NCBI Protein Information	angiopoietin-1 receptor; hTIE2; p140 TEK; soluble TIE2 variant 1; soluble TIE2 variant 2; endothelial tyrosine kinase; tyrosine-protein kinase receptor TEK; tunica interna endothelial cell kinase; tyrosine-protein kinase receptor TIE-2; tyrosine kinase wi
UniProt Protein Name	Angiopoietin-1 receptor
UniProt Gene Name	TEK
UniProt Synonym Gene Names	TIE2; VMCM; VMCM1; hTIE2
UniProt Entry Name	TIE2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.